



Cochrane
Library

Cochrane Database of Systematic Reviews

Remote versus face-to-face check-ups for asthma (Review)

Kew KM, Cates CJ

Kew KM, Cates CJ.

Remote versus face-to-face check-ups for asthma.

Cochrane Database of Systematic Reviews 2016, Issue 4. Art. No.: CD011715.

DOI: 10.1002/14651858.CD011715.pub2.

www.cochranelibrary.com

Remote versus face-to-face check-ups for asthma (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	7
OBJECTIVES	8
METHODS	8
RESULTS	10
Figure 1.	11
Figure 2.	14
DISCUSSION	17
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	20
CHARACTERISTICS OF STUDIES	28
DATA AND ANALYSES	45
Analysis 1.1. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 1 Exacerbations requiring oral corticosteroids.	46
Analysis 1.2. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 2 Exacerbations requiring hospital emergency department (ED) treatment.	47
Analysis 1.3. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 3 Exacerbations requiring hospital admission.	48
Analysis 1.4. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 4 Asthma control (Asthma Control Questionnaire (ACQ)).	49
Analysis 1.5. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 5 Serious adverse events (including mortality).	49
Analysis 1.6. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 6 Asthma-related quality of life (Asthma Quality of Life Questionnaire (AQLQ)).	50
Analysis 1.7. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 7 Unscheduled healthcare visits.	51
Analysis 1.8. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 8 Change in lung function (trough FEV1).	52
Analysis 1.9. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 9 Adverse events.	52
Analysis 2.1. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 1 Exacerbations requiring hospital admission.	53
Analysis 2.2. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 2 Asthma control (ACQ).	53
Analysis 2.3. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 3 Asthma-related quality of life (AQLQ).	54
Analysis 2.4. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 4 Unscheduled healthcare visits.	54
Analysis 2.5. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 5 Adverse events.	55
ADDITIONAL TABLES	55
APPENDICES	56
CONTRIBUTIONS OF AUTHORS	59
DECLARATIONS OF INTEREST	59
SOURCES OF SUPPORT	59
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	59

[Intervention Review]

Remote versus face-to-face check-ups for asthma

Kayleigh M Kew¹, Christopher J Cates¹

¹Population Health Research Institute, St George's, University of London, London, UK

Contact address: Kayleigh M Kew, Population Health Research Institute, St George's, University of London, Cranmer Terrace, London, SW17 0RE, UK. kkew@sgul.ac.uk.

Editorial group: Cochrane Airways Group.

Publication status and date: New, published in Issue 4, 2016.

Review content assessed as up-to-date: 24 November 2015.

Citation: Kew KM, Cates CJ. Remote versus face-to-face check-ups for asthma. *Cochrane Database of Systematic Reviews* 2016, Issue 4. Art. No.: CD011715. DOI: 10.1002/14651858.CD011715.pub2.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Asthma remains a significant cause of avoidable morbidity and mortality. Regular check-ups with a healthcare professional are essential to monitor symptoms and adjust medication.

Health services worldwide are considering telephone and internet technologies as a way to manage the rising number of people with asthma and other long-term health conditions. This may serve to improve health and reduce the burden on emergency and inpatient services. Remote check-ups may represent an unobtrusive and efficient way of maintaining contact with patients, but it is uncertain whether conducting check-ups in this way is effective or whether it may have unexpected negative consequences.

Objectives

To assess the safety and efficacy of conducting asthma check-ups remotely versus usual face-to-face consultations.

Search methods

We identified trials from the Cochrane Airways Review Group Specialised Register (CAGR) up to 24 November 2015. We also searched www.clinicaltrials.gov, the World Health Organization (WHO) trials portal, reference lists of other reviews and contacted trial authors for additional information.

Selection criteria

We included parallel randomised controlled trials (RCTs) of adults or children with asthma that compared remote check-ups conducted using any form of technology versus standard face-to-face consultations. We excluded studies that used automated telehealth interventions that did not include personalised contact with a health professional. We included studies reported as full-text articles, as abstracts only and unpublished data.

Data collection and analysis

Two review authors screened the literature search results and independently extracted risk of bias and numerical data. We resolved any disagreements by consensus, and we contacted study authors for missing information.

We analysed dichotomous data as odds ratios (ORs) using study participants as the unit of analysis, and continuous data as mean differences using the random-effects models. We rated all outcomes using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.

Remote versus face-to-face check-ups for asthma (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

1

Main results

Six studies including a total of 2100 participants met the inclusion criteria: we pooled four studies including 792 people in the main efficacy analyses, and presented the results of a cluster implementation study (n = 1213) and an oral steroid tapering study (n = 95) separately. Baseline characteristics relating to asthma severity were variable, but studies generally recruited people with asthma taking regular medications and excluded those with COPD or severe asthma. One study compared the two types of check-up for oral steroid tapering in severe refractory asthma and we assessed it as a separate question. The studies could not be blinded and dropout was high in four of the six studies, which may have biased the results.

We could not say whether more people who had a remote check-up needed oral corticosteroids for an asthma exacerbation than those who were seen face-to-face because the confidence intervals (CIs) were very wide (OR 1.74, 95% CI 0.41 to 7.44; 278 participants; one study; low quality evidence). In the face-to-face check-up groups, 21 participants out of 1000 had exacerbations that required oral steroids over three months, compared to 36 (95% CI nine to 139) out of 1000 for the remote check-up group. Exacerbations that needed treatment in the Emergency Department (ED), hospital admission or an unscheduled healthcare visit all happened too infrequently to detect whether remote check-ups are a safe alternative to face-to-face consultations. Serious adverse events were not reported separately from the exacerbation outcomes.

There was no difference in asthma control measured by the Asthma Control Questionnaire (ACQ) or in quality of life measured on the Asthma Quality of Life Questionnaire (AQLQ) between remote and face-to-face check-ups. We could rule out significant harm of remote check-ups for these outcomes but we were less confident because these outcomes are more prone to bias from lack of blinding.

The larger implementation study that compared two general practice populations demonstrated that offering telephone check-ups and proactively phoning participants increased the number of people with asthma who received a review. However, we do not know whether the additional participants who had a telephone check-up subsequently benefited in asthma outcomes.

Authors' conclusions

Current randomised evidence does not demonstrate any important differences between face-to-face and remote asthma check-ups in terms of exacerbations, asthma control or quality of life. There is insufficient information to rule out differences in efficacy, or to say whether or not remote asthma check-ups are a safe alternative to being seen face-to-face.

PLAIN LANGUAGE SUMMARY

Are telephone or internet check-ups a safe alternative to being seen face-to-face?

Take-home message

Studies that tried to answer this research question did not show important differences between the two types of check-up. However, there is not enough information to rule out differences in their harms or benefits. At this stage, we cannot say whether or not asthma check-ups conducted over the phone or internet are a safe alternative to usual face-to-face consultations.

Background

Regular contact with a doctor or asthma nurse is essential to keep track of symptoms and use of inhalers. Telephone and internet technologies may be a way to manage the rising number of people with asthma and other long-term health conditions. This has been referred to as 'remote reviews' or e-consultations, and may be a way of more easily keeping contact between patients and doctors, but we don't know whether it's as good as meeting face-to-face.

Study characteristics

We found a total of six studies including 2100 participants: four studies including 792 people could be pooled for the main results, and two other studies were looked at separately because their designs were very different (n = 1213 and n = 95). People in the four pooled studies in general took regular medications and we excluded those with severe asthma or other lung diseases. We looked at two other studies with very different designs to the main four separately: one compared a practice where people with asthma were given the option of a telephone check-up or a practice visit where they came to the clinic as usual, and one looked specifically at using technology to monitor people while cutting down their oral steroids dose. We last looked for studies on 24 November 2015.

Key results

We cannot say whether or not people who had a check-up over the phone or internet were more or less likely to need oral corticosteroids for an asthma attack than those seen face-to-face, and we were uncertain of the result for several reasons. Too few people had asthma attacks that needed treatment in the Emergency Department or hospital, or an unscheduled visit to see their doctor to tell if remote check-ups were as good as face-to-face consultations. There didn't appear to be a difference in asthma control or quality of life, but we were able to rule out the possibility that remote check-ups are not as good as face-to-face consultations on these measures. The evidence was all considered to be of low or moderate quality. The study that tested the possible benefit of giving people the option of a telephone check-up showed that this increased the number of people reviewed, but did not show an overall benefit on asthma outcomes.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Remote versus face-to-face check-ups for asthma							
Patient or population: adults or children with asthma Setting: outpatient Intervention: remote check-ups conducted using technology (e.g. telephone, email) Comparison: face-to-face asthma check-ups							
Outcomes	Anticipated absolute effects* (95% CI)			Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with face-to-face check-ups	Risk with check-ups	remote check-ups				
Exacerbations requiring oral corticosteroids 3 months	21 per 1000	36 per 1000 (9 to 139)		OR 1.74 (0.41 to 7.44)	278 (1 RCT)	⊕⊕○○ low ^{1,2}	Very imprecise. Data from the implementation study** were consistent
Exacerbations requiring hospital admission 6 months	5 per 1000	3 per 1000 (0 to 33)		Peto OR 0.63 (0.06 to 6.32)	651 (3 RCTs)	⊕⊕○○ low ^{1,3}	Very few events - no conclusion could be drawn. The implementation study was more in favour of face-to-face check-ups
Asthma control (ACQ) Scale 0 to 6; lower is better 12 months	The mean ACQ score with face-to-face check-ups improved by 0.11	The mean ACQ score with remote check-ups improved by 0.07 more (0.35 more to 0.21 less)	-		146 (1 RCT)	⊕⊕⊕○ moderate ^{4,5}	No difference and CIs ruled out significant harm of remote check-ups (MCID for the ACQ is 0.5). The implementation study results were consistent

Serious adverse events (including mortality)	-	-	-	0 RCTs	-	No efficacy studies reported all-cause SAEs. The implementation study recorded 12/554 and 8/659 in the remote and face-to-face groups respectively (OR 1.80, 95% CI 0.73 to 4.44)
Asthma-related quality of life (AQLQ) Scale 1 to 7; higher is better 8 months	The mean AQLQ score with face-to-face check-ups was 5.49	The mean AQLQ score with remote check-ups was 0.08 better (0.14 worse to 0.30 better)	-	544 (3 RCTs)	⊕⊕⊕○ moderate ^{4,5}	No difference and CIs ruled out significant harm of remote check-ups (MCID for the AQLQ is 0.5). The implementation study results were consistent
Unscheduled health-care visits 5 months	120 per 1000	110 per 1000 (58 to 201)	OR 0.91 (0.45 to 1.85)	531 (2 RCTs)	⊕⊕○○ low ^{1,3}	Very few events - we could not draw any conclusions. The implementation study was more precise and did not show a difference
Lung function (trough FEV ₁) 6 months	The mean trough FEV ₁ with face-to-face check-ups was 20 mL	The mean trough FEV ₁ with remote check-ups was 166.76 mL better (78.03 more to 255.5 more)	-	253 (1 RCT)	⊕⊕⊕○ moderate ^{1,6}	People having remote check-ups had better lung function than those seen face-to-face in the one study that measured it

***The risk in the intervention group** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI). The risk with face-to-face check-ups for continuous outcomes was calculated as a weighted mean of the face-to-face values.

Abbreviations: CI = confidence interval; RR = risk ratio; OR = odds ratio; ED = emergency department; ACQ = Asthma Control Questionnaire; AQLQ = Asthma Quality of Life Questionnaire; FEV₁ = forced expiratory volume in one second; RCT = randomised controlled trial

**The 'Implementation study', [Pinnock 2007a](#), had a two-cluster pragmatic design and was not pooled with the rest of the included studies (efficacy studies)

Durations were calculated as a weighted mean duration of the studies contributing data to the analysis

GRADE Working Group grades of evidence

High quality: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate quality: we are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low quality: our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect.

Very low quality: we have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹Studies were at high risk of bias for one or more blinding domains but it is unlikely that this had an effect on the objective outcomes (no downgrade).

²Evidence from 1 study with 7 events. There were very wide CIs (downgrade by 2 for imprecision).

³The effect was based on very few events. The 95% confidence intervals included significant harm and significant benefit of remote check-ups (downgraded by 2 for imprecision).

⁴The upper limit of the CI crossed the line of no effect but both limits were well within the 0.5 unit minimal clinically important difference for the scale (no downgrade).

⁵Studies were at high risk of bias for blinding which may have affected this subjective outcome, and there was evidence of possible attrition bias (downgraded by 1 for risk of bias).

⁶The CIs excluded benefit of face-to-face check-ups but they were wide and based on only one study of 253 people (downgraded by 1 for imprecision).

BACKGROUND

Description of the condition

Asthma is a chronic disease of the airways, which causes reversible inflammation and narrowing of the airways and mucus production (GINA 2014). It commonly causes symptoms of wheezing, breathlessness, chest tightness and cough, although these vary between people and over time in their presence, frequency and severity (GINA 2014).

Despite the emergence and update of several national and international management guidelines which recommend a range of cost-effective treatments based on frequency and severity of symptoms and exacerbations (e.g. BTS/SIGN 2014; GINA 2014), the disease remains a significant cause of avoidable morbidity and mortality worldwide (BTS/SIGN 2014; Global Asthma Report 2014; NRAD 2014). A national review of the 195 asthma deaths that occurred between February 2012 and January 2013 in the UK revealed that, in the year preceding their death, nearly one-third had no record of seeing a general practitioner (GP) and nearly two-thirds had not had an asthma check-up in secondary care (NRAD 2014). The importance of self-monitoring and regular check-ups with a healthcare professional to monitor symptoms, and encourage adherence to prevent inhalers, is now well accepted (Gibson 2002; NRAD 2014), especially for people at high risk of severe asthma attacks.

Description of the intervention

Communication technologies, such as telephones and video conferencing, have been proposed as a way to conduct asthma check-ups remotely. Conducting check-ups in this way is a form of 'telehealth', otherwise referred to as 'telecare', 'digital health', 'telemedicine' or 'e-health'. McLean 2013 described this field as "the use of information and communication technologies to deliver healthcare at a distance and to support patient self-management through remote monitoring and personalised feedback". It may also be conceptualised as "an emerging field in the intersection of medical informatics, public health and business, referring to health services and information delivered or enhanced through the internet and related technologies (Eysenback 2001). Health services around the world are considering remote check-ups as a way to manage the rising number of people with long-term health conditions, to improve health outcomes and reduce the burden on emergency and inpatient services (Department of Health 2012; Steventon 2012).

The UK government outlined its aims for the widespread use of technology in health in their 2013 mandate, including wide availability of 'e-consultations' by GPs, and significant progress towards home 'telemonitoring' of three million people with long-term conditions by 2017 (Department of Health 2013). Researchers have

studied the role of a range of technology-based check-ups and monitoring in asthma and other health conditions, including the use of telephone calls, email contact, text-messaging and video-conferencing (Laver 2013; McLean 2010; McLean 2011).

How the intervention might work

In the context of asthma, a condition that affects around 334 million people worldwide (Global Asthma Report 2014) and places a significant burden on healthcare systems, remote check-ups may represent an unobtrusive and efficient way of maintaining contact with patients. Regular monitoring with communication technologies that does not disrupt a patient's life in the way that regular clinic visits might, may serve to enhance self management behaviours that have known benefits on morbidity and mortality, such as keeping personalised action plans up-to-date, and adherence to maintenance medications (NRAD 2014).

However, while governments and health services have highlighted the potential for cost savings and improved clinical outcomes of using remote check-ups instead of face-to-face consultations, its use to monitor patients with potentially serious or life-threatening conditions may not be without hazard. Focus groups have suggested that telehealth may be acceptable to patients and clinicians, but they have also raised concerns that it could actually discourage self management, or increase the likelihood of serious outcomes, by instilling a false sense of security (Pinnock 2007b).

The feasibility of using communication technologies in different situations and populations may be hampered by barriers, including insufficient healthcare infrastructure and funding (Lustig 2012). However, it may be a way to reduce inequality in health care related to socioeconomic status and rural living by improving access to services (Jannett 2003; Lustig 2012).

Why it is important to do this review

The release of the UK National Health Service (NHS) mandate in 2013 has seen a push to advance the use of telehealth for economic and clinical benefit. A recent overview of systematic reviews suggested that these benefits should not be assumed and that people at highest risk of serious health outcomes are likely to show the biggest gains (McLean 2013). For asthma, existing reviews have noted a large degree of variation in the way telehealth is defined and delivered in studies, to whom and to what it is compared (Jaana 2009; McLean 2010), and have been limited for this reason in the conclusions that could be drawn. This Cochrane review will focus on conducting asthma check-ups remotely as a form of telehealth compared with usual face-to-face consultations in a hospital or clinic. A related Cochrane review will consider the evidence for remote monitoring of asthma control between visits with ongoing personalised feedback from a health professional (Kew 2015a).

OBJECTIVES

To assess the safety and efficacy of conducting asthma check-ups remotely versus usual face-to-face consultations.

METHODS

Criteria for considering studies for this review

Types of studies

We included parallel randomised controlled trials (RCTs) of any duration. We included studies reported as full-text articles, those published as an abstract only and unpublished data.

Types of participants

We included studies of adults or children with a diagnosis of asthma. We excluded studies that recruited participants with other long-term health conditions, unless they presented data for people with asthma separately.

Types of interventions

We included trials that compared remote check-ups conducted with any form of technology (e.g. telephone calls, video-conferencing) versus standard face-to-face check-ups. We included trials which compared the two types of check-up on top of education or another co-intervention. We excluded trials that used automated telehealth interventions and did not include personalised contact with a health professional.

Types of outcome measures

Primary outcomes

1. Exacerbations that required oral corticosteroids^a.
2. Asthma control (measured on a validated scale, e.g. the Asthma Control Questionnaire (ACQ)).
3. Serious adverse events (including mortality).

^aIf trials reported exacerbations in a different way (e.g. required hospital emergency department (ED) visit), we analysed these separately.

Secondary outcomes

1. Asthma-related quality of life (measured on a validated scale, e.g. the Asthma Quality of Life Questionnaire (AQLQ)).
2. Unscheduled healthcare visits.
3. Lung function (trough forced expiratory volume in one second (FEV₁) preferred).

4. Adverse events/side effects.

Reporting of one or more of the outcomes listed here in the trial was not an inclusion criterion for this Cochrane review.

Search methods for identification of studies

Electronic searches

We identified trials from the Cochrane Airways Review Group's Specialised Register (CAGR), which is maintained by the Information Specialist for the Cochrane Airways Review Group. The CAGR contains trial reports identified through systematic searches of bibliographic databases including the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, EMBASE, CINAHL, AMED, and PsycINFO, and handsearches of respiratory journals and meeting abstracts (please see [Appendix 1](#) for further details). We used the search strategy in [Appendix 2](#) to search for all records in the CAGR.

We also conducted a search of ClinicalTrials.gov (www.ClinicalTrials.gov) and the World Health Organization (WHO) International Clinical Trials Registry Portal (www.who.int/ictrp/en/) limited to interventional studies, using the condition term 'asthma', intervention terms 'remote OR telehealth OR telemedicine OR internet OR web', and title terms 'NOT education'. We searched all databases from their inception to 24 November 2015, and did not impose any restriction on language of publication.

Searching other resources

We checked reference lists of all primary studies and review articles for additional references.

We searched for errata or retractions from included studies published in full-text on PubMed (www.ncbi.nlm.nih.gov/pubmed) on 20 July 2015.

Data collection and analysis

Selection of studies

Two review authors (KK and CJC) independently screened titles and abstracts for inclusion of all the potential studies identified from the literature searches and coded them as either 'retrieve' (eligible or potentially eligible/unclear) or 'do not retrieve'. We retrieved the full-text study reports/publication and two review authors (KK and CJC) independently screened the full-text and identified studies for inclusion, and identified and recorded reasons for exclusion of the ineligible studies. We resolved any disagreement through discussion. We identified and excluded duplicates and collated multiple reports of the same study so that each

study, rather than each report, was the unit of interest in the review. We recorded the selection process in sufficient detail to complete a PRISMA flow diagram and 'Characteristics of excluded studies' table (Moher 2009).

Data extraction and management

We used a data collection form for study characteristics and outcome data, which we piloted on at least one study included in the review. One review author (KK) extracted the following study characteristics from included studies.

1. Methods: study design, total duration of study, details of any 'run in' period, number of study centres and location, study setting, withdrawals and date of study.
2. Participants: number of participants, mean age, age range, gender, severity of condition, diagnostic criteria, baseline lung function, smoking history, inclusion criteria and exclusion criteria.
3. Interventions: intervention, comparison, concomitant medications and excluded medications.
4. Outcomes: primary and secondary outcomes specified and collected, and time points reported.
5. Notes: trial funding and notable conflicts of interest of the trial authors.

Two review authors (KK and CJC) independently extracted outcome data from the included studies. We noted in the 'Characteristics of included studies' table if the study authors did not report outcome data in a usable way. We resolved any disagreements by consensus. One review author (KK) transferred data into the Review Manager (RevMan) (RevMan 2014) file. We double-checked that KK entered data correctly by comparing the data presented in the systematic review with the study reports. A second review author (CJC) spot-checked study characteristics for accuracy against the trial report.

Assessment of risk of bias in included studies

Two review authors (KK and CJC) independently assessed the risk of bias for each included study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We resolved any disagreements by discussion. We assessed the risk of bias according to the following domains.

1. Random sequence generation.
2. Allocation concealment.
3. Blinding of participants and personnel.
4. Blinding of outcome assessment.
5. Incomplete outcome data.
6. Selective outcome reporting.
7. Other bias.

We graded each potential source of bias as either 'high', 'low' or 'unclear', and provided a quote from the study report together with a justification for our judgment in the 'Risk of bias' table. We summarised the 'Risk of bias' judgements across different studies

for each domain listed. We considered blinding separately for different key outcomes where necessary (e.g. for unblinded outcome assessment, risk of bias for all-cause mortality may be very different than for a patient-reported pain scale). Where information on risk of bias related to unpublished data or correspondence with a trial author, we noted this in the 'Risk of bias' table.

When we considered treatment effects, we took into account the risk of bias for the studies that contributed to that outcome.

Assessment of risk of bias in conducting the systematic review

We conducted the review according to the published Cochrane protocol, Kew 2015b, and reported any deviations from it in the 'Differences between protocol and review' section.

Measures of treatment effect

We analysed dichotomous data as odds ratios (ORs) and continuous data as either mean difference or standardised mean difference values. We entered data presented as a scale with a consistent direction of effect.

We undertook meta-analyses only where this was meaningful, i.e. if the treatments, participants and the underlying clinical question were similar enough for pooling to make sense.

We narratively described skewed data reported as medians and interquartile ranges.

Where a single trial reported multiple trial arms, we only included the relevant trial arms. If the trial combined two comparisons (e.g. drug A versus placebo and drug B versus placebo) in the same meta-analysis, we halved the control group to avoid double-counting.

Unit of analysis issues

For dichotomous outcomes we used participants, rather than events, as the unit of analysis (i.e. number of adults admitted to hospital, rather than number of admissions per adult). However, if studies reported exacerbations as rate ratios, we analysed them on this basis. We did not anticipate the inclusion of cluster RCTs and hence presented a large two-cluster implementation study, Pinnock 2007a, separately from the other studies. For the purposes of display in the analyses, we have referred to Pinnock 2007a as the 'cluster implementation study' and Chan 2007, Gruffydd-Jones 2005, Pinnock 2003 and Rasmussen 2005 as the 'efficacy RCTs'. There were only two clusters so we included the data with participants as the unit of analysis. Although we presented the results of the cluster RCT on forest plots with the other studies, we did not pool the effects so weighting of the cluster trial within the analysis was not an issue. We presented Hashimoto 2011 in a separate comparison as the study focused on tapering OCS dose.

Dealing with missing data

We contacted the study authors or study sponsors in order to verify key study characteristics and obtain missing numerical outcome data where possible (e.g. when we identified a study as an abstract only). Where this was not possible, and we thought the missing data introduced serious bias, we explored the impact of including such studies in the overall assessment of results by a sensitivity analysis.

Assessment of heterogeneity

We used the I^2 statistic to measure heterogeneity among the trials in each analysis. If we identified substantial heterogeneity we reported it and explored possible causes by prespecified subgroup analysis.

Assessment of reporting biases

If we were able to pool more than 10 trials, we created and examine a funnel plot to explore possible small study and publication biases.

Data synthesis

We used a random-effects model for all analyses, as we expected variation in effects due to differences in study populations and interventions. We performed sensitivity analyses with a fixed-effect model when heterogeneity was statistically significant.

'Summary of findings' table

We created a 'Summary of findings' table using the seven outcomes listed above. We used the five GRADE considerations (study limitations, consistency of effect, imprecision, indirectness and publication bias) to assess the quality of a body of evidence as it relates to the studies which contribute data to the meta-analyses for the prespecified outcomes. We used methods and recommendations described in Section 8.5 and Chapter 12 of the *Cochrane Handbook for Systematic Reviews of Interventions*, Higgins 2011, and used GRADEpro Guidelines Development Tool (GDT) software (GRADEpro GDT 2015). We justified all decisions to downgrade or upgrade the quality of the evidence in footnotes, and we made comments to aid the reader's understanding of the Cochrane review, where necessary.

Subgroup analysis and investigation of heterogeneity

We performed the following subgroup analyses for the primary outcomes, where there was a sufficient number of included studies.

1. Mean age (less than 16 years, 17 to 65 years, and greater than 65 years).
2. Type of technology (telephone calls, text-messages, emails).

We used the formal test for subgroup interactions in Review Manager (RevMan) (RevMan 2014).

Sensitivity analysis

We carried out the following sensitivity analyses by exclusion of the following from the primary analyses.

1. Studies that recruited people with severe or life-threatening asthma.
2. Unpublished data (obtained from trial authors or from conference abstracts).
3. Studies at high risk of selection bias^a.

^aInadequate selection procedures may result in unbalanced baseline characteristics between groups which could skew the data. Due to the nature of the studies, we anticipated that all or most included studies would be at high risk of performance or detection bias, so we discussed the possible effect of lack of blinding, in particular for subjective outcomes.


RESULTS

Description of studies

Results of the search

We performed searches up to 24 November 2015. We identified 591 records from the Cochrane Airways Review Group's Specialised Register (CAGR). We also examined a total of 750 additional records, comprised of an older database search (n = 710), clinicaltrials.gov records (n = 29) and the World Health Organization (WHO) International Clinical Trials Registry Portal (ICTRP) (n = 11). After we removed duplicates, we screened the remaining 823 records and excluded 673 by looking at titles and abstracts alone. We retrieved full texts for the remaining 150 records and excluded 126 (collated into 95 studies). The other 25 records met all the inclusion criteria and we collated them as six included studies. We have presented a study flow diagram and the reasons for exclusion in Figure 1.

Figure 1. Study flow diagram



Included studies

Six studies including a total of 2100 participants met the inclusion criteria: we pooled four studies including 792 people in the main efficacy analyses, and presented the results of a cluster implementation study (n = 1213) and an oral steroid tapering study (n = 95) separately. We have presented a summary of study, participant and intervention characteristics in [Table 1](#), and more details for each individual study are in the 'Characteristics of included studies' tables.

Design, setting and duration

Three included studies were conducted in the UK ([Gruffydd-Jones 2005](#); [Pinnock 2003](#); [Pinnock 2007a](#)), one in Hawaii, USA ([Chan 2007](#)), one in the Netherlands ([Hashimoto 2011](#)) and one in Denmark ([Rasmussen 2005](#)). Five studies were randomised controlled trials (RCTs) that lasted six or 12 months, and [Pinnock 2007a](#) was a before-and-after implementation study which randomised three practices rather than individual participants.

The design of [Pinnock 2007a](#) differed from the other studies in several respects. Firstly, it was cluster randomised which had an effect on the unit used for analysis and could not completely control for potential group differences between the practices. Secondly, it was a before-and-after design which meant the number and type of people on the practice lists was different at the two time points. Thirdly, the intervention offered a telephone check-up as an option for asthma review so the study did not make a clean comparison between remote and face-to-face check-ups, especially since most people in the telephone group did not choose that option. Fourthly, the primary aim of the study was to increase the number of people having a check-up at all and so people in the intervention group were phoned opportunistically on top of being offered a phone check-up, and this additional effort to contact participants may have been a confounding factor on the other study outcomes. We considered the study to be important so did not exclude it, and instead presented the study results separately from the main comparison for each outcome.

Population characteristics and inclusion criteria

The number of participants in each trial ranged from 95 in [Hashimoto 2011](#) to 1728 in [Pinnock 2007a](#); the median number of participants recruited was 236 and the total who received phone or face-to-face check-ups was 2100. This was complicated by the design of [Pinnock 2007a](#) which assigned three GP practices rather than individual participants, although we will discuss the number of people for the descriptive purposes. Five studies recruited adults with a lower age limit of 17 or 18 and an upper age limit of between 45 ([Rasmussen 2005](#)) and 75 ([Hashimoto 2011](#)).

Mean age of participants in the adult trials ranged from 29 to 55.5 years (median 50.1). One study recruited children between the ages of six and 17 years, and had a mean age of 9.6 years ([Chan 2007](#)). Studies included slightly more females than males (range 34.5% to 62.5% male, median 45%). Baseline characteristics related to the asthma severity were patchy and variable across studies. [Hashimoto 2011](#) recruited people with more severe asthma than the other included studies; the mean percentage predicted FEV₁ at baseline was 73.9%. Two other studies reported this measure of baseline severity at 92% ([Rasmussen 2005](#)) and 100.5% ([Chan 2007](#)).

In general the included studies did not describe the inclusion and exclusion criteria in great detail, and had varying requirements for the diagnosis and classification of asthma in their participants. With the exception of [Hashimoto 2011](#), the studies recruited from the practice or clinic lists of their participating centres. [Pinnock 2003](#) and [Pinnock 2007a](#) required participants to have received a prescription for asthma medications within the previous six and 12 months respectively, and both excluded participants with chronic obstructive disease. [Pinnock 2003](#) further required participants to have been diagnosed with asthma for at least a year. Other inclusion criteria were related to computer or telephone access, and exclusion criteria were related to social, communication or medical difficulties that might preclude involvement in the intervention. [Hashimoto 2011](#) only recruited participants on daily oral corticosteroids, and its inclusion criteria differed from the other studies. This study required participants to have a diagnosis of severe refractory asthma according to the major and minor criteria recommended by the American Thoracic Society ([ATS 2000](#)), and for their asthma to be uncontrolled despite intensive follow-up by an asthma specialist for at least a year, chronic treatment with oral corticosteroids and high doses of inhaled steroids and long-acting bronchodilators.

Interventions and comparisons

The interventions received in the active and comparison groups varied in several respects across studies, in particular regarding the length of observation, amount of professional contact and method of communication.

Four studies were designed to test remote check-ups with standard face-to-face care. [Chan 2007](#) was the only child study, and tested an in-home website case management and education programme against an in-person equivalent for 12 months. Participants in the active group of [Gruffydd-Jones 2005](#) received phone calls every six months from trained asthma nurses and those in the intervention group had equivalent face-to-face check-ups with the asthma nurse. Both groups were given a personalised asthma action plan and discussed symptoms, peak flow and inhaler tech-

nique. [Pinnock 2003](#) was described as a pragmatic RCT where participants were offered a telephone check-up or a face-to-face consultation in the surgery, both with an asthma nurse. One of the study's main aims was the uptake of check-ups with either method within three months of randomisation. [Rasmussen 2005](#) was a six-month study where physicians gave participants instructions via email or over the phone based on an agreed asthma action plan, an online diary and peak flow measurements uploaded by the participant. The comparison group received face-to-face instruction from an asthma specialist on how to adjust medication based on their asthma action plan and peak flow measurements.

As described above, [Pinnock 2007a](#) was a 12-month implementation study where participants were given a structured recall with a choice of telephone or face-to-face for their asthma check-up or a structured recall with no choice of a telephone check-up (i.e. face-to-face only). We did not include a usual-care group not subject to the methods to control for bias in the main comparison in this review. We chose to present the results of this study alongside the results of the four studies above, but in a separate subgroup so we did not pool the data with the main comparison. In addition to the choice of a telephone check-up, people in the intervention group were called opportunistically if they did not respond or attend, which did not happen in the control group.

[Hashimoto 2011](#) compared face-to-face check-ups with internet-delivered check-ups for the specific purpose of tapering long-term oral corticosteroid therapy (OCS), and for this reason we analysed the study as a separate comparison. The study was designed to test the effectiveness of a six-month programme of OCS adjustment either via an internet diary and associated monitoring from

an asthma nurse or via face-to-face check-ups with a specialist according to Global Initiative for Asthma (GINA) guidelines.

Excluded studies

We excluded 125 records after viewing full texts, which we collated to represent 94 unique studies. The most common reason for exclusion was that the intervention did not meet the inclusion criteria (82 studies). Within this explanation, we explored the reasons and found that we often excluded studies because they used technology to facilitate self-monitoring between usual face-to-face check-ups (n = 22), and these studies meet the inclusion for another Cochrane review ([Kew 2015a](#)). Other excluded studies with interventions that did not meet the inclusion criteria used technology as education rather than for asthma check-ups (n = 14), as automated monitoring systems without involvement from a health professional (n = 8), to validate an asthma questionnaire (n = 7), to improve or monitor adherence (n = 7), or to deliver a range of other interventions that did not match the remit of this review (including technology delivered counselling or behavioural interventions, parenting advice and monitoring interventions delivered without technology). We excluded eight studies as they were not RCTs (n = 8), and four because they made the wrong comparison (n = 4).

Risk of bias in included studies

We have shown a summary of the 'Risk of bias' judgements in [Figure 2](#).

Figure 2. 'Risk of bias' summary: review authors' judgements about each 'Risk of bias' item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel: Objective outcomes	Blinding of participants and personnel: Subjective outcomes	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Chan 2007	+	?	+	-	?	-	+	+
Gruffydd-Jones 2005	+	?	+	-	-	-	+	+
Hashimoto 2011	+	?	+	-	-	-	+	+
Pinnock 2003	+	+	+	-	?	+	+	+
Pinnock 2007a	+	-	+	-	?	+	+	-
Rasmussen 2005	+	+	+	-	-	-	+	+

Allocation

We rated all the included studies at low risk of bias for random sequence generation as they described the randomisation methods, such as centralised systems, random numbers tables or coin toss. We judged two studies at low risk of bias for allocation concealment after the study authors responded to our request to clarify the study methods. Also we considered one study at unclear risk of bias because it did not adequately describe the methods. Regarding [Pinnock 2007a](#), the cluster implementation study, we rated it at high risk of bias for allocation concealment because of its two-cluster design.

Blinding

Due to the nature of the interventions, none of the included studies were able to blind participants and personnel to group allocation (performance bias). For this reason, we chose to assess blinding of participants and personnel separately for the subjective and objective outcomes. In each study, we rated the subjective outcomes (Asthma Control Questionnaire (ACQ) and Asthma Quality of Life Questionnaire (AQLQ)) at high risk of bias and the objective outcomes (exacerbations, adverse events, FEV₁) at low risk of bias. While it was possible for studies to blind outcome assessors (detection bias), no included study described the procedures to do so. [Gruffydd-Jones 2005](#), [Hashimoto 2011](#) (through personal communication) and [Rasmussen 2005](#) explicitly stated that the outcomes assessors were not blinded to allocation so we rated them at high risk of bias, and the remaining studies as unclear.

Incomplete outcome data

We considered two included studies at low risk of bias ([Pinnock 2003](#); [Pinnock 2007a](#)). The former had low and even dropout, and the latter was a real-world implementation study where uptake rate was an integral part of the study. We rated four studies at high risk of bias due to incomplete outcome data, either because dropout was high or unbalanced between groups or missing data had not been sufficiently imputed to account for those not in the study at the end (or both) ([Chan 2007](#); [Gruffydd-Jones 2005](#); [Hashimoto 2011](#); [Rasmussen 2005](#)). Due to the nature of the question being posed in the studies, several were run in a real-world context which made it more difficult to control for participants dropping out.

Selective reporting

There was no evidence of selective reporting in the six included studies so we rated them at low risk of bias. Where outcomes were unavailable in the published reports, the trial authors were able to provide the additional data or confirm they had not been measured.

Other potential sources of bias

We rated [Pinnock 2007a](#) at high risk of bias for several reasons related to its cluster before-and-after study design. It randomised two practices to the interventions which would not have controlled for baseline imbalances in the same way as individual randomisation, and this meant the participant population in each group was not static. The intervention was a telephone option and many in that practice opted for a usual face-to-face check-up, which meant the study did not make a direct comparison of remote and face-to-face check-ups. Additionally, people in the telephone option group were phoned opportunistically to increase uptake of check-ups, which did not happen in the face-to-face group. These factors mean we cannot be certain that the mode of check-up, and not the increased likelihood of being seen, was the variable measured and we considered this study as having a high risk of bias. For these reasons, we chose to present the study alongside the others but not to pool its results in the main analyses.

We did not identify any other sources of bias in the five other included studies, which we rated as having a low risk of bias.

Effects of interventions

See: [Summary of findings for the main comparison Summary of findings table](#)

We have presented data from studies that made a direct comparison between remote and face-to-face check-ups as the main results (referred herein as the 'efficacy studies', and supplemented these by the results from the large two-cluster implementation study, [Pinnock 2007a](#). We have described data from the OCS tapering study, [Hashimoto 2011](#), as a separate comparison below.

Primary outcomes

Exacerbations that required oral corticosteroids

One efficacy study reported the number of people who needed a course of oral corticosteroids for an exacerbation of asthma ([Pinnock 2003](#)). The confidence intervals (CIs) were very wide due to the small number of events in the analysis (odds ratio (OR) 1.74, 95% CI 0.41 to 7.44; 278 participants). In the face-to-face check-up group, 21 people out of 1000 had exacerbations requiring oral steroids over three months, compared to 36 (95% CI nine to 139) out of 1000 in the remote group. We downgraded the evidence to low quality due to imprecision. The effect from the cluster implementation study was much more precise and favoured face-to-face check-ups (OR 1.43, 95% CI 1.04 to 1.97); we have shown the two effects together in [Analysis 1.1](#).

Exacerbations that required emergency department (ED) visit and hospital admission

The effect for exacerbations that required treatment in the ED also favoured face-to-face check-ups over those conducted remotely, but this was uncertain due to the wide CIs from a small number of events (OR 2.60, 95% CI 0.63 to 10.64; 651 participants; three studies; [Analysis 1.2](#)). The result of the cluster implementation study was much smaller but also very imprecise (OR 1.19, 95% CI 0.38 to 3.71; 1212 participants; one study). Too few people in the efficacy studies had exacerbations that required a hospital admission to detect any difference between the two types of check-up (Peto OR 0.63, 95% CI 0.06 to 6.32; 651 participants; three studies; [Analysis 1.3](#)), and the cluster implementation study showed a possible but not statistically significant benefit of face-to-face check-up (Peto OR 2.18, 95% CI 0.83 to 5.69; 1213 participants; one study).

Asthma control

There was no difference in scores on the Asthma Control Questionnaire (ACQ) between participants in the remote and face-to-face groups (mean difference (MD) -0.07 , 95% CI -0.35 to 0.21 ; 146 participants; one study; [Analysis 1.4](#)). Both CIs were within the minimal clinically important difference (MCID) for the scale (MCID = 0.5). We downgraded the quality of the evidence to moderate quality due to risk of bias because the outcome was a subjective rating scale that may have been affected by the inability to blind participants and personnel to group allocation. While the effect from the cluster implementation study was more in favour of remote check-ups than the efficacy studies, the estimate and its CIs were still within the MCID for the scale so the difference was unimportant.

Serious adverse events (including mortality)

Only the cluster implementation study reported serious adverse events (SAEs) and showed a higher number in participants in the remote groups, although the CIs did not exclude the possibility that they were more common with face-to-face check-ups (OR 1.80, 95% CI 0.73 to 4.44; 1213 participants; one study; [Analysis 1.5](#)). Given that SAEs are usually defined as those requiring hospital admission, it is likely that most of these events were the exacerbations requiring hospital admission described above.

Subgroup analyses

We were unable to conduct meaningful subgroup analyses on the basis of age and type of technology as planned, as there was an insufficient number of included studies to do this.

Sensitivity analyses

Severe or life-threatening asthma

It was not necessary to conduct this sensitivity analysis because only [Hashimoto 2011](#) specifically recruited people with severe or life-threatening asthma and we analysed this study on its own due to the nature of the intervention.

Unpublished data

The trial author of [Pinnock 2007a](#) provided additional data for exacerbations requiring oral steroids, but we did not pool the study effect with the other study in the analysis so there was no basis for a sensitivity analysis.

High risk of selection bias

None of the included studies were at high risk of selection biases so we did not perform the planned sensitivity analysis.

Secondary outcomes

Asthma-related quality of life

There was no difference between remote and face-to-face check-ups on the Asthma Quality of Life Questionnaire (AQLQ) (MD 0.08, 95% CI -0.14 to 0.30 ; 544 participants; three studies; [Analysis 1.6](#)). While the effect was marginally in favour of remote check-ups, both CIs were within the 0.5 minimal clinically important difference on the scale. As with the ACQ, we downgraded the evidence to moderate quality for risk of bias because the outcome was subjective and may have been affected by lack of blinding. The point estimate in the implementation study lay marginally in the opposite direction but the result was not inconsistent given the MCID (-0.02 , 95% CI -0.23 to 0.19 ; 536 participants; one study).

[Rasmussen 2005](#) also reported the AQLQ but the data were skewed and analysed non-parametrically so we could not combine it with the other studies in the meta-analysis (internet group median 6.42, range 4.11 to 7.00; specialist group median 6.17, range 3.98 to 7.00; GP group median 6.31, range 1.41 to 7.00). [Chan 2007](#) also reported the parent version of the AQLQ with the following scores (remote group: mean 6.4, SD = 1, N = 60; face-to-face group: mean 6.2, SD = 0.8, N = 60). We considered the child version to be more similar to the way the scores were taken in the other studies which is why we included the child and not the parent scores in the meta-analysis.

Unscheduled healthcare visits

The pooled estimate from two efficacy studies was based on very few events and was too imprecise to draw a conclusion (OR 0.91, 95% CI 0.45 to 1.85; 531 participants; two studies; [Analysis 1.7](#)). The direction and magnitude of the effect from the implementation study were consistent but the estimate was more precise (OR 0.95, 95% CI 0.75 to 1.21; 1213 participants; one study).

Lung function (trough FEV₁ preferred)

Only one efficacy study reported trough FEV₁ (mL), and showed a bigger improvement in the remote check-up group (MD 166.76, 95% CI 78.03 to 255.50; 253 participants; one study; [Analysis 1.8](#)). We downgraded the evidence for imprecision to moderate quality because it was based on only 253 people, even though the confidence intervals excluded the possibility that face-to-face check-ups were better. The implementation study was not designed to measure lung function.

Adverse events/side effects

Studies generally did not report adverse events separately from the asthma exacerbation and resource use outcomes. One study author confirmed that no participants in either group experienced adverse events, which did not allow us to display an effect estimate.

Remote versus face-to-face check-up for oral corticosteroid tapering - analysis of Hashimoto 2011

We chose to assess [Hashimoto 2011](#) separately from the other studies as it was specifically aimed at assessing remote versus face-to-face check-ups to guide dose reduction for people with asthma taking long-term oral corticosteroids.

Mostly the CIs were too wide to tell whether one type of check-up was better than the other at reducing harmful outcomes that might occur as a result of withdrawing oral steroids, or to say that they were equivalent. This was true for exacerbations requiring hospital admission (OR 0.88, 95% CI 0.25 to 3.13; [Analysis 2.1](#)), unscheduled healthcare visits (OR 2.31, 95% CI 0.23 to 23.14; [Analysis 2.4](#)) and adverse events (OR 1.51, 95% CI 0.13 to 17.29; [Analysis 2.5](#)). Scores on the ACQ (MD 0.14, 95% CI -0.15 to 0.43; [Analysis 2.2](#)) and AQLQ (MD -0.17, 95% CI -0.49 to 0.15; [Analysis 2.3](#)) were slightly better in the face-to-face group than those who were managed remotely, but the difference was below the minimal clinically important difference on both scales and the CIs did not exclude the possibility that remote check-ups were better.

DISCUSSION

Summary of main results

Six studies met the inclusion criteria (including a total of 2100 participants): we pooled four studies in the main efficacy analyses, which randomised 792 people to remote or face-to-face check-up. In addition we also presented the results of a cluster implementation study (n = 1213) alongside the main results but did not pool them with the other included studies, and assessed one study that compared the two types of check-up for oral steroid tapering in severe refractory asthma as a separate comparison (n = 95). Baseline characteristics relating to asthma severity were variable, but studies generally recruited people with asthma who took regular medications and excluded those with chronic obstructive pulmonary disease (COPD) or severe asthma. The studies could not be blinded and dropout was high in four of the six included studies, which may have biased the results.

We cannot say whether more people in the remote groups needed oral corticosteroids for an asthma exacerbation than those seen face-to-face because the confidence intervals (CIs) were very wide (OR 1.74, 95% CI 0.41 to 7.44; 278 participants; one study; low quality evidence). In the face-to-face check-up groups, 21 people out of 1000 had exacerbations that required oral steroids over three months, compared to 36 (95% CI nine to 139) out of 1000 for the remote check-up group. Exacerbations that needed treatment in the hospital emergency department (ED), hospital admission or an unscheduled healthcare visit happened too infrequently to detect whether remote check-ups are a safe alternative to face-to-face consultations. Serious adverse events were not reported separately from the exacerbation outcomes.

There was no difference in asthma control measured by the Asthma Control Questionnaire (ACQ) or in quality of life measured on the Asthma Quality of Life Questionnaire (AQLQ) between remote and face-to-face check-ups. We ruled out the significant harm of remote check-ups for these outcomes but we were less confident because these outcomes are more prone to bias from lack of blinding.

The larger implementation study that compared two general practice populations demonstrated that offering telephone check-ups and proactively phoning participants increased the number of people with asthma who received a review. However, we do not know whether the additional participants who had a telephone check-up subsequently benefited in asthma outcomes.

Overall completeness and applicability of evidence

The design of this Cochrane review allowed us to focus on the specific question of whether regular asthma check-ups can effectively be conducted remotely as opposed to face-to-face. By honing in on this form of telehealth, we were able to assess the evidence for a clearly defined application of technology-based care that was more likely to lead to evidence that could be readily applied to real world

settings. However, it did limit our assessments of possible effect moderators, such as age and type of technology, due to the small number of studies that met the inclusion criteria and reported the primary outcomes.

As described in the 'Included studies' section, [Pinnock 2007a](#) was a real-world implementation study that differed from the other included studies in several respects. We chose to include this study because it addressed the feasibility of remote check-ups in a real-world setting, despite its design being dissimilar to the other included studies. By not pooling it with the other studies but presenting the results alongside them, we hoped to highlight the possible differences in a controlled comparison of remote and face-to-face check-ups and what might actually happen in practice. While [Pinnock 2007a](#)'s results do not lead to the cause and effect inferences that can be made from the efficacy RCTs, it does give an important insight into the relative merits of a possible incorporation of remote check-ups into practice, and we consider the review to be more complete as a result. We did not set out to assess the possible benefit of improving access by offering remote check-ups, especially for patients who are less likely to attend their annual check-up. This is an important factor that can be addressed by implementation studies, such as [Pinnock 2007a](#), better than classic randomised controlled trials (RCTs). Unfortunately, because there were only two clusters, the study results must be interpreted with caution and with the caveats both in this Cochrane review and in the study itself.

One efficacy study contributed data to the exacerbations requiring oral steroids analysis and this was provided by the study author ([Pinnock 2003](#)). It was not always possible to distinguish between courses of oral steroids for immediate treatment or future use (as part of an action plan). This highlights the importance of contacting study authors for additional data, as their own primary outcomes may differ from those of the review. This increased the completeness of the evidence base and our confidence in the results.

Quality of the evidence

We rated the evidence in this Cochrane review as low or moderate quality, mostly due to imprecision or risk of bias. We chose to assess blinding for the subjective outcomes separately from the objective outcomes. By differentiating the outcomes in this way, we were able to assess the effect lack of blinding was likely to have had on the confidence we have in the results. For the objective outcomes, we noted that the study designs did not allow for blinding of participants and personnel, but judged that this lack of blinding is unlikely to have made a difference to the number of people who had exacerbations, adverse events and lung function. There is a possibility that the lack of blinding may have still affected the way these outcomes were recorded but we did not consider this to be sufficient to downgrade the evidence. For the subjective outcomes, it is more likely that the participant or investigators'

knowledge of group allocation could have affected the way they responded to the asthma control and quality of life questionnaires, so we downgraded these outcomes for risk of bias.

We downgraded five outcomes for imprecision, including the primary outcome. We were fairly certain in the direction of the effect for the primary outcome, exacerbations requiring oral steroids, but there was only one efficacy study in the analysis and seven events, which reduced our confidence in the size and precision of the estimate significantly. Exacerbations that required an ED visit or a hospital admission, and unscheduled healthcare visits were all based on a very small number of events, which crucially meant we could not rule out significant harm of conducting remote check-ups for people with asthma. Finally, we downgraded trough FEV₁ for imprecision for a similar reason to the primary outcome: the effect was in favour of remote check-ups, but the estimate was based on just one study of 253 people, so we weren't confident in the result. Visually, there was imprecision in the estimates for asthma control and quality of life, but the CIs for both were well within the established minimal clinically important differences for the scales ([Juniper 1994](#); [Juniper 1999](#)).

We did not downgrade any of the outcomes for publication bias, inconsistency in the results or indirectness of the evidence. While not all studies reported the outcomes we were interested in, this was more likely to reflect the individual practices and designs of the studies (e.g. unable to measure lung function remotely) rather than selective reporting of outcome data. There was minimal statistical inconsistency in the analyses, and while there was a fair amount of variation in the aims and designs of the studies, we considered them all to match the PICO set out in the protocol for this review ([Kew 2015b](#)).

Potential biases in the review process

We made every effort to adhere to Cochrane methods during the review process. Both review authors extracted numerical data, performed 'Risk of bias' ratings in duplicate, and cross checked for accuracy and consistency. Throughout the process, we resolved any discrepancies through discussion. Neither review author has any conflicts of interest relating to this Cochrane review.

We performed broad literature searches that we independently screened in duplicate, and included studies regardless of language of publication or the existence of a full-text paper. We also conducted comprehensive additional searches to identify unpublished studies that were not listed in the main electronic databases. It is unlikely that we missed studies during the study selection process, except for those not listed in non-English language databases. We made every effort to contact study authors, although in some cases this was difficult due to when the studies were conducted and the effect this had both on the availability of contact details and the likelihood we could obtain data. We received detailed replies from two study authors which affected several analyses and the completeness of the 'Risk of bias' information.

Agreements and disagreements with other studies or reviews

Systematic reviews of the evidence for remote asthma check-ups have generally had much broader inclusion criteria, assessing the use of any kind of telehealthcare for monitoring or educating people with asthma. As such, it is difficult to compare the results of this Cochrane review with their results because we designed this systematic review to assess specifically the use of technology systems to conduct asthma check-ups remotely. For example, [McLean 2010](#) included 21 studies that assessed a range of telehealth interventions against a range of control groups, and as such it was difficult to meta-analyse the data in a way that led to conclusions with real-world implications.

Systematic reviews that examined telehealth more broadly for asthma have generally approached the question from a different standpoint from our own. Other reviews have focused on possible benefits of remote healthcare over usual face-to-face care, whereas we were more concerned with highlighting the potential dangers of removing face-to-face contact with a healthcare professional. Conclusions made by [McLean 2010](#) regarding the lack of benefit for relatively mild asthma and [Zhao 2014](#) regarding the lack of improvements on asthma function scores do not disagree with our own, but are framed from the former standpoint. This is at least partly because these two reviews did not set out to compare remote or 'telehealthcare' against a consistent alternative form of monitoring. For most outcomes we looked at, we cannot conclude that remote check-ups definitely lead to worse outcomes than face-to-face care, but neither can we conclude that remote check-ups are a safe alternative to face-to-face care for either mild or severe asthma.

In their conclusions, [Jaana 2009](#) focused on the attitudes and receptiveness of home telemonitoring for patients with respiratory illness, which we did not set out to assess, and commented on the "variations in study approaches and an absence of robust study designs and formal evaluations". We agree that the inherent differences in health systems and study designs continue to limit conclusions in this field, even when review questions are refined to one part of 'telehealthcare'.

AUTHORS' CONCLUSIONS

Implications for practice

Current randomised evidence does not demonstrate any important differences between face-to-face and remote asthma check-ups in terms of exacerbations, asthma control or quality of life. There is insufficient information to rule out differences in efficacy, or to say whether or not remote asthma check-ups are a safe alternative to usual face-to-face consultations.

Implications for research

It would be helpful to find out whether remote check-ups for people with asthma who do not attend for a face-to-face consultation can achieve a reduction in the risk of serious asthma attacks that result in hospital admissions. To do this, further studies are required in high-risk people with frequent asthma admissions who do not attend for regular face-to-face check-up. It might be sensible to include remote monitoring and remote check-ups in the intervention as it seems unlikely that remote check-ups alone would be sufficient to reduce admissions to hospital.

ACKNOWLEDGEMENTS

We are grateful to the Cochrane Airways Group (CAG) staff for comments and support. We thank Simone Hashimoto, Hilary Pinnock and Linda Makowska Rasmussen who provided additional data and clarified the methods for three of the included studies.

Rebecca Normansell was the Editor for this review and commented critically on this review.

The [Background](#) and [Methods](#) sections of this review are based on a standard template used by Cochrane Airways.

The [Background](#) section shares similarities with another Cochrane review we co-developed ([Kew 2015a](#)).

The National Clinical Guideline Centre (NCGC) and the CAG undertook collaborative work pertaining to a systematic review of published evidence on tele-healthcare for monitoring asthma control. The CAG reviews are restricted to interventions that involve a healthcare professional only. This is different from the larger question addressed by the NCGC (as part of the National Institute for Health and Care Excellence (NICE) asthma guideline commission). The NCGC review of evidence is published in the NICE clinical guideline on asthma diagnosis and monitoring and received funding from NICE.

REFERENCES

References to studies included in this review

Chan 2007 *{published and unpublished data}*

Callahan CW. Asthma in-home monitoring (AIM) trial. <http://www.clinicaltrials.gov/ct/show/NCT00282516> (accessed 14 August 2015).

Callahan CW, Chan DS, Hatch-Pigott G, Manning N, Proffitt L, Lawless A, et al. One year randomized controlled trial of home telemonitoring of children with persistent asthma using store-and-forward technology vs office-based care [Abstract]. American Thoracic Society 2005 International Conference; May 20-25; San Diego, California. 2005:[C47] [Poster: A21].

* Chan DS, Callahan CW, Hatch-Pigott VB, Lawless A, Proffitt HL, Manning NE, et al. Internet-based home monitoring and education of children with asthma is comparable to ideal office-based care: results of a 1-year asthma in-home monitoring trial. *Pediatrics* 2007;**119**(3): 569–78.

Chan DS, Callahan CW, Sheets SJ, Moreno CN, Malone FJ. An Internet-based store-and-forward video home telehealth system for improving asthma outcomes in children. *American Journal of Health-System Pharmacy* 2003; **60**(19):1976–81.

Gruffydd-Jones 2005 *{published and unpublished data}*

* Gruffydd-Jones K, Hollinghurst S, Ward S, Taylor G. Targeted routine asthma care in general practice using telephone triage. *British Journal of General Practice* 2005;**55** (521):918–23.

Gruffydd-Jones K, Ward S. Targeted routine asthma care in general practice using the telephone triage [Abstract]. *European Respiratory Journal*. 2005; Vol. 26:Abstract No. 4264.

Hashimoto 2011 *{published and unpublished data}*

Hashimoto S, Ten Brinke A, Roldaan AC, van Veen IH, Moller GM, Sont JK, et al. Monitoring exhaled Nitric Oxide (FENO) To Tailor The Lowest Effective Dose Of Oral Corticosteroids In Severe Asthma (MONOSA-Study) [Abstract]. *American Journal of Respiratory and Critical Care Medicine*. 2010; Vol. 181:A3721.

* Hashimoto S, Ten Brinke A, Roldaan AC, van Veen IH, Möller GM, Sont JK, et al. Internet-based tapering of oral corticosteroids in severe asthma: A pragmatic randomised controlled trial. *Thorax* 2011;**66**(6):514–20.

Pinnock 2003 *{published and unpublished data}*

* Pinnock H, Bawden R, Proctor S, Wolfe S, Scullion J, Price D, et al. Accessibility, acceptability, and effectiveness in primary care of routine telephone review of asthma: pragmatic, randomised controlled trial. *BMJ* 2003;**326** (7387):477–9.

Pinnock H, Sheikh A, Bawden R, Proctor S, Wolfe S, Scullion J, et al. A randomised controlled trial comparing telephone review with fact to face consultations in the

management of adult asthmatics in UK primary care.

Primary Care Respiratory Journal 2002;**11**:68–9.

Pinnock H, Sheikh A, Bawden R, Proctor S, Wolfe S, Scullion J, et al. Cost effectiveness of telephone vs face to face consultations for annual asthma review: randomised controlled trial in UK primary care [Abstract]. *European Respiratory Journal*. 2003; Vol. 22:Abstract No. 2250.

Pinnock H, Sheikh A, Bawden R, Proctor S, Wolfe S, Scullion J, et al. Telephone vs. face to face consultations in the management of adult asthmatics in UK primary care: a randomised controlled trial [Abstract]. *European Respiratory Society 12th Annual Congress*; 2002 Sep 14–18; Stockholm. 2002:Abstract No. P593.

Pinnock 2007a *{published and unpublished data}*

* Pinnock H, Adlem L, Gaskin S, Harris J, Snellgrove C, Sheikh A. Accessibility, clinical effectiveness, and practice costs of providing a telephone option for routine asthma reviews: phase IV controlled implementation study. *British Journal of General Practice* 2007;**57**(542):714–22.

Pinnock H, Adlem L, Gaskin S, Snellgrove C, Harris J, Sheikh A. Impact on access of providing a telephone option for primary care asthma reviews: controlled implementation study [Abstract]. *European Respiratory Journal*. 2005; Vol. 26:Abstract No. 1711.

Pinnock H, Madden V, Snellgrove C, Sheikh A. Telephone or surgery asthma reviews? Preferences of participants in a primary care randomised controlled trial. *Primary Care Respiratory Journal* 2005;**14**:42–6.

Pinnock H, McKenzie L, Price D, Sheikh A. Cost-effectiveness of telephone or surgery asthma reviews: economic analysis of a randomised controlled trial. *British Journal of General Practice* 2005;**55**(511):119–24.

Pinnock H, Norman C, Bowden K, Sheikh A. Impact on asthma morbidity and patient enablement of providing a telephone option for primary care asthma reviews: phase IV controlled implementation study [Abstract]. *Primary Care Respiratory Journal*. 2006; Vol. 15:187.

Pinnock H, Norman C, Bowden K, Sheikh A. Impact on asthma morbidity and patient enablement of providing a telephone option in a primary care asthma review service controlled implementation study [Abstract]. *European Respiratory Journal*. 2006; Vol. 28:571s [3361].

Rasmussen 2005 *{published and unpublished data}*

Phanareth K, Rasmussen L, Nolte H, Backer V. Using the Internet as a tool for the management of asthma disease. *European Respiratory Journal* 2002;**20**:54s.

Rasmussen L M, Phanareth K, Nolte H, Backer V. Can internet-based monitoring of asthma maintain asthma control over a 12 month period [Abstract]. *European Respiratory Journal*. 2006; Vol. 28:122s [783].

Rasmussen L, Backer V, Phanareth K. Preliminary data from a clinical trial: does use of an internet based asthma-monitoring tool increase lung function and asthma control? [Abstract]. *American Thoracic Society 99th International*

Conference; 2003 May 16-21; Seattle. 2003:A102 Poster D6.

Rasmussen L, Phanareth K, Nolte H, Backer V. A long term randomized clinical study of 300 asthmatics. An internet based asthma monitoring tool improves quality of life significantly [Abstract]. *European Respiratory Journal*. 2004; Vol. 24:261s.

Rasmussen L, Phanareth K, Nolte H, Backer V. Preliminary data from a clinical trial: An Internet based asthma monitoring tool increases lung function and decrease airway hyperresponsiveness [Abstract]. *European Respiratory Journal*. 2003; Vol. 22:Abstract No: [3351].

Rasmussen LM, Phanareth K, Nolte H, Backer V. Can internet-based management improve asthma control? A long term randomised clinical study of 300 asthmatics [Abstract]. *European Respiratory Journal*. 2005; Vol. 26: Abstract No. 400.

* Rasmussen LM, Phanareth K, Nolte H, Backer V. Internet-based monitoring of asthma: a long-term, randomized clinical study of 300 asthmatic subjects. *Journal of Allergy and Clinical Immunology* 2005;**115**(6):1137–42.

References to studies excluded from this review

ACTRN12606000400561 {published data only}

ACTRN12606000400561. Improving childhood asthma management through a telemedicine monitoring network. <https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?ACTRN=12606000400561> (accessed 2 June 2015).

Xu C, Jackson M, Scuffham PA, Wootton R, Simpson P, Whitty J, et al. A randomized controlled trial of an interactive voice response telephone system and specialist nurse support for childhood asthma management. *Journal of Asthma* 2010;**47**(7):768–73.

Ahmed 2011 {published data only}

Ahmed S, Bartlett S J, Ernst P, Pare G, Kanter M, Perreault R, et al. Effect of a web-based chronic disease management system on asthma control and health-related quality of life: Study protocol for a randomized controlled trial. *Trials* 2011;**12**:260.

Ahmed S, Bartlett SJ, Ernst P, Lin CJ, Pare G, Perreault R, et al. My asthma portal: preliminary results of a web-based self management intervention [Abstract]. *American Journal of Respiratory and Critical Care Medicine*. 2011; Vol. 183: A5321.

Andersen 2007 {published data only}

Andersen UM. Does a www-based interactive computer program change asthma outcomes, quality of life and asthma knowledge [Abstract]. *Journal of Allergy and Clinical Immunology*. 2007; Vol. 119:S9 [34].

Apter 2000 {published data only}

Apter A. Inhaled steroid adherence in moderate and severe asthma. CRISP (Computer Retrieval of Information on Scientific Projects) 2000; Vol. 31 August 2005.

Apter 2015 {published data only}

Apter AJ, Bryant-Stephens, Morales KH, Wan F, Hardy S, Reed-Wells S, et al. Using IT to improve access, communication, and asthma in African American and Hispanic/Latino Adults: Rationale, design, and methods of a randomized controlled trial. *Contemporary Clinical Trials* 2015;**44**:119–28.

Araujo 2012 {published data only}

Araujo L, Jacinto T, Moreira A, Castel-Branco M G, Delgado L, Costa-Pereira A, et al. Clinical efficacy of web-based versus standard asthma self-management. *Journal of Investigational Allergology and Clinical Immunology* 2012;**22**(1):28–34.

Baptist 2013 {published data only}

Baptist AP, Ross JA, Yang Y, Song PX, Clark NM. A randomized controlled trial of a self-regulation intervention for older adults with asthma. *Journal of the American Geriatrics Society* 2013;**61**(5):747–53.

Barbanel 2003 {published data only}

Barbanel D, Eldridge S, Griffiths C. Can a self-management programme delivered by a community pharmacist improve asthma control? A randomised trial. *Thorax* 2003;**58**(10): 851–4.

Bateman 2000 {published data only}

Bateman ED, Kruger MJ. A computer-based home-monitoring disease management programme, pulmassist plus® (PAP), achieves significant improvement in quality of life, and healthcare costs in moderate and severe asthma [Abstract]. *American Journal of Respiratory and Critical Care Medicine*. 2000; Vol. 161:A457.

Bender 2001 {published data only}

Bender BG. Assessment mode and validity of self-reports in children. CRISP (Computer Retrieval of Information on Scientific Projects) 2001; Vol. 31 January 2005 End date.

Bender 2007 {published data only}

Bender BG, Bartlett SJ, Rand CS, Turner C, Wamboldt FS, Zhang L. Impact of interview mode on accuracy of child and parent report of adherence with asthma-controller medication. *Pediatrics* 2007;**120**(3):e471–7.

Bender 2010 {published data only}

Bender BG, Apter A, Bogen DK, Dickinson P, Fisher L, Wamboldt FS, et al. Test of an interactive voice response intervention to improve adherence to controller medications in adults with asthma. *Journal of the American Board of Family Medicine* 2010;**23**(2):159–65.

Boyd 2014 {published data only}

Boyd MJ, Elliott RA, Barber N, Mehta R, Waring J, Chuter A, et al. The impact of the New Medicines Service (NMS) in England on patients adherence to their medicines. *International Journal of Pharmacy Practice* 2014;**22**(S2):66.

Burbank 2012 {published data only}

Burbank A, Rettiganti M, Brown RH, Jones S, Perry TT. Asthma education via telemedicine: Effects on asthma knowledge and self-efficacy [Abstract]. *Journal of Investigative Medicine*. 2012; Vol. 60:401.

- Burkhardt 2002** *{published data only}*
Burkhardt P. Promoting Children's Adherence to Asthma Self-Management. CRISP (Computer Retrieval of Information on Scientific Projects) 2002; Vol. 31 July 2004 End date.
- Bynum 2001** *{published data only}*
Bynum A, Hopkins D, Thomas A, Copeland N, Irwin C. The effect of telepharmacy counseling on metered-dose inhaler technique among adolescents with asthma in rural Arkansas. *Telemedicine Journal and E-Health* 2001;7(3):207–17.
- Chandler 1990** *{published data only}*
Chandler MH, Clifton GD, Louis BA, Coons SJ, Foster TS, Phillips BA. Home monitoring of theophylline levels: a novel therapeutic approach. *Pharmacotherapy* 1990;10(4):294–300.
- Chatkin 2006** *{published data only}*
Chatkin JM, Blanco DC, Scaglia N, Wagner MB, Fritscher CC. Impact of a low-cost and simple intervention in enhancing treatment adherence in a Brazilian asthma sample. *Journal of Asthma* 2006;43(4):263–6.
- Chen 2013** *{published data only}*
Chen SH, Huang JL, Yeh KW, Tsai YF. Interactive support interventions for caregivers of asthmatic children. *Journal of Asthma* 2013;50(6):649–57.
- Cicutto 2009** *{published data only}*
Cicutto L, Ashby MN, Feldman D. Telephone intervention based strategies to increase the completion and use of asthma action plans for adults with asthma [Abstract]. American Thoracic Society International Conference; 2009 May 15-20; San Diego. 2009:A1033.
- Clark 2007** *{published data only}*
Clark NM, Gong ZM, Wang SJ, Lin X, Bria WF, Johnson TR. A randomized trial of a self-regulation intervention for women with asthma. *Chest* 2007;132(1):88–97.
- Clarke 2014** *{published data only}*
Clarke SA, Calam R, Morawska A, Sanders M. Developing web-based Triple P 'Positive Parenting Programme' for families of children with asthma. *Child: Care, Health and Development* 2014;40(4):492–7.
- Claus 2004** *{published data only}*
Claus R, Michael H, Josef L, Jan-Torsten T, Marion S. Internet based patient education evaluation of a new tool for young asthmatics [Abstract]. European Respiratory Society 14th Annual Congress; 2004 Sep 3-7; Glasgow. 2004; Vol. 24:383s.
- Cruz-Correia 2007** *{published data only}*
Cruz-Correia R, Fonseca J, Lima L, Araujo L, Delgado L, Castel-Branco MG, et al. A comparison of web-based and paper-based self management tools for asthma: Patients' opinions and quality of data in a randomised crossover study. *Journal on Information Technology in Healthcare* 2007; 5:357–71.
Cruz-Correia R, Fonseca J, Lima L, Araújo L, Delgado L, Castel-Branco MG, et al. Web-based or paper-based self-management tools for asthma--patients' opinions and quality of data in a randomized crossover study. *Studies in Health Technology and Informatics* 2007;127:178–89.
- de Jongste 2008** *{published data only}*
de Jongste JC, Carroro S, Hop WC, Baraldi E. Daily exhaled nitric oxide telemonitoring in the management of childhood asthma [Abstract]. American Thoracic Society International Conference; 2008 May 16-21; Toronto. 2008:Poster #908.
- Deschildre 2012** *{published data only}*
Deschildre A, Béghin L, Salleron J, Iliescu C, Thumerelle C, Santos C, et al. Home telemonitoring (forced expiratory volume in 1 s) in children with severe asthma does not reduce exacerbations. *European Respiratory Journal* 2012;39(2):290–6.
- De Vera 2014** *{published data only}*
De Vera MA, Sadatsafavi M, Tsao NW, Lynd LD, Lester R, Gastonguay L, et al. Empowering pharmacists in asthma management through interactive SMS (EmPhAsIS): study protocol for a randomized controlled trial. *Trials* 2014;15:488.
- Donald 2008** *{published data only}*
Donald KJ, McBurney H, Teichtahl H, Irving L. A pilot study of telephone based asthma management. *Australian Family Physician* 2008;37(3):170–3.
- Dwinger 2013** *{published data only}*
Dwinger S, Dirmaier J, Herbarth L, König HH, Eckardt M, Kriston L, et al. Telephone-based health coaching for chronically ill patients: study protocol for a randomized controlled trial. *Trials* 2013;14:337.
- Eakin 2012** *{published data only}*
Eakin MN, Rand CS, Bilderback A, Bollinger ME, Butz A, Kandasamy V, et al. Asthma in Head Start children: effects of the Breathmobile program and family communication on asthma outcomes. *Journal of Allergy and Clinical Immunology* 2012;129(3):664–70.
- Finkelstein 2005** *{published data only}*
Finkelstein J. Inhaled steroid adherence in moderate & severe asthma. CRISP (Computer Retrieval of Information on Scientific Projects) 2000; Vol. 31 July 2005 End date.
* Finkelstein J, Joshi A, Amelung P. Evaluation of home telemanagement in adult asthma patients [Abstract]. *Journal of Allergy and Clinical Immunology* 2005;115(2):S63.
Finkelstein J, Joshi A, Arora M, Amelung P. Impact of home telemanagement in adult asthma [Abstract]. American Thoracic Society 2005 International Conference; May 20-25; San Diego. 2005:A40; Poster: G14.
- Fonseca 2006** *{published data only}*
Fonseca JA, Costa-Pereira A, Delgado L, Fernandes L, Castel-Branco MG. Asthma patients are willing to use mobile and web technologies to support self-management. *Allergy* 2006;61(3):389–90.
- Foster 2014** *{published data only}*
Foster J, Smith L, Usherwood T, Sawyer S, Reddel H. Electronic reminders improve adherence with preventer inhalers in Australian primary care patients [Abstract].

- Annual Scientific Meetings of the Thoracic Society of Australia and New Zealand and the Australian and New Zealand Society of Respiratory Science; 2014 April 4-9; Adelaide. 2014:24–34.
- Friedman 1999** *{published data only}*
Friedman R. Impact of a telecommunication system in childhood asthma. CRISP (Computer Retrieval of Information on Scientific Projects) 1999; Vol. 29 September 2004 End date.
- Garbutt 2010** *{published data only}*
* Garbutt JM, Banister C, Highstein G, Sterkel R, Epstein J, Bruns J, et al. Telephone coaching for parents of children with asthma: impact and lessons learned. *Archives of Pediatrics and Adolescent Medicine* 2010;**164**(7):625–30.
Swerczek LM, Banister C, Bloomberg GR, Bruns JM, Epstein J, Highstein GR, et al. A telephone coaching intervention to improve asthma self-management behaviors. *Pediatric Nursing* 2013;**39**(3):125–130, 145.
- Guendelman 2002** *{published data only}*
* Guendelman S, Meade K, Benson M, Chen YQ, Samuels S. Improving asthma outcomes and self-management behaviors of inner-city children: a randomized trial of the Health Buddy interactive device and an asthma diary. *Archives of Pediatrics and Adolescent Medicine* 2002;**156**(2): 114–20.
Guendelman S, Meade K, Chen YQ, Benson M. Asthma control and hospitalizations among inner-city children: results of a randomized trial. *Telemedicine Journal and e-Health* 2004;**10**(Suppl 2):S–6–14.
- Gustafson 2012** *{published data only}*
* Gustafson D, Wise M, Bhattacharya A, Pulvermacher A, Shanovich K, Phillips B, et al. The effects of combining web-based eHealth with telephone nurse case management for pediatric asthma control: A randomized controlled trial. *Journal of Medical Internet Research* 2012;**14**(4):e101.
Wise M, Gustafson DH, Sorkness CA, Molfenter T, Staresinic A, Meis T, et al. Internet telehealth for pediatric asthma case management: integrating computerized and case manager features for tailoring a web-based asthma education program. *Health Promotion Practice* 2007;**8**(3): 282–91.
- Halterman 2012** *{published data only}*
* Halterman JS, Fagnano M, Montes G, Fisher S, Tremblay P, Tajon R, et al. The school-based preventive asthma care trial: results of a pilot study. *Journal of Pediatrics* 2012;**161**(6):1109–1115.e1.
Halterman JS, Sauer J, Fagnano M, Montes Gu, Fisher S, Tremblay P, et al. Working toward a sustainable system of asthma care: development of the School-Based Preventive Asthma Care Technology (SB-PACT) trial. *Journal of Asthma* 2012;**49**(4):395–400.
- Huang 2013** *{published data only}*
Huang JL, Chen SH, Yeh KW. Health outcomes, education, healthcare delivery and quality-3056. Constructed supporting program improves asthma treatment outcomes in children. *World Allergy Organization Journal* 2013;**6**(suppl 1):P224.
- Jan 2007** *{published data only}*
Jan RL, Wang JY, Huang MC, Tseng SM, Su HJ, Liu LF, et al. An internet-based interactive tele-monitoring system for improving childhood asthma outcomes in Taiwan. *Telemedicine Journal and e-Health* 2007;**13**(3):257–68.
- Janevic 2012** *{published data only}*
Janevic MR, Sanders GM, Thomas LJ, Williams DM, Nelson B, Gilchrist E, et al. Study protocol for Women of Color and Asthma Control: a randomized controlled trial of an asthma-management intervention for African American women. *BMC Public Health* 2012;**12**:76.
- Jerant 2003** *{published data only}*
Jerant A. A Randomized Trial of Home Self-Efficacy Enhancement. CRISP (Computer Retrieval of Information on Scientific Projects) 2003; Vol. 31 December 2007 End date.
- Kattan 2006** *{published data only}*
Kattan M, Crain EF, Steinbach S, Visness CM, Walter M, Stout JW, et al. A randomized clinical trial of clinician feedback to improve quality of care for inner-city children with asthma. *Pediatrics* 2006;**117**(6):e1095–103.
- Khan 2003** *{published data only}*
Khan MSR, O'Meara M, Stevermuier TL, Henry RL. Randomized controlled trial of asthma education after discharge from an emergency department. *Journal of Paediatrics and Child Health* 2004;**40**(12):674–7.
* Khan S, O'Meara M, Hurst T, Henry RL. Randomised controlled trial of asthma education by telephone after discharge from an emergency department [Abstract]. *European Respiratory Journal* 2003;**22**:Abstract No: [P2294].
- Kojima 2005** *{published data only}*
Kojima N, Takeda Y, Akashi M, Kamiya T, Matsumoto M, Ohya Y, et al. Interactive education during summer camp for children with asthma improved adherence of self-management [Abstract]. *Journal of Allergy and Clinical Immunology* 2005;**115**:S115.
- Kokubu 1999** *{published data only}*
Kokubu F, Nakajima S, Ito K, Makino S, Kitamura S, Fukuchi Y, et al. [Hospitalization reduction by an asthma tele-medicine system]. *Aerugi [Allergy]* 2000;**49**(1):19–31.
* Kokubu F, Suzuki H, Sano Y, Kihara N, Adachi M. [Tele-medicine system for high-risk asthmatic patients]. *Aerugi [Allergy]* 1999;**48**(7):700–12.
- Lam 2011** *{published data only}*
Lam A. Practice innovations: delivering medication therapy management services via Videoconference interviews. *Consultant Pharmacist* 2011;**26**(10):764–74.
- Liu 2011** *{published data only}*
* Liu WT, Huang CD, Wang CH, Lee KY, Lin SM, Kuo HP. A mobile telephone-based interactive self-care system improves asthma control. *European Respiratory Journal* 2011;**37**(2):310–7.
Liu WT, Wang CH, Huang CD, Kuo HP, Liu WT, Kuo HP. A novel mobile phone-based self-care system improves asthma control [Abstract]. American Thoracic Society

- International Conference; 2007 May 18-23; San Francisco. 2007:[A93].
- Lobach 2013** *{published data only}*
Lobach DF, Kawamoto K, Anstrom KJ, Silvey GM, Willis JM, Johnson FS, et al. A randomized trial of population-based clinical decision support to manage health and resource use for Medicaid beneficiaries. *Journal of Medical Systems* 2013;**37**(1):9922.
- McCowan 2001** *{published data only}*
McCowan C, Neville RG, Ricketts IW, Warner FC, Hoskins G, Thomas GE. Computer assisted assessment and management of patients with asthma. American Thoracic Society 97th International Conference; 2001 May 18-23; San Francisco. 2001.
- McPherson 2006** *{published data only}*
Glazebrook C, McPherson A, Forster D, James C, Crook I, Smyth A. The asthma files: randomised controlled trial of interactive multimedia program to promote self-management skills in children [Abstract]. American Thoracic Society 100th International Conference; 2004 May 21-26; Orlando. 2004:D92 Poster 506.
* McPherson AC, Glazebrook C, Forster D, James C, Smyth A. A randomized, controlled trial of an interactive educational computer package for children with asthma. *Pediatrics* 2006;**117**(4):1046–54.
- Merchant 2013** *{published data only}*
Merchant R, Inamdar R, Quade R, Van Sickle D, Maenner M, Patmas M. Interim results from a randomized, controlled trial of remote monitoring of inhaled bronchodilator use on asthma control and management [Abstract]. American College of Chest Physicians Annual Meeting; 26-31 October 2013; Chicago, Illinois. 2013; Vol. 144:71A.
- Morrison 2014** *{published data only}*
Morrison D, Wyke S, Thomson NC, McConnachie A, Agur K, Saunderson K, et al. A Randomized trial of an Asthma Internet Self-management Intervention (RAISIN): study protocol for a randomized controlled trial. *Trials* 2014;**15**:185.
- Murphy 2001** *{published data only}*
Murphy JC. Telemedicine offers new way to manage asthma. *American Journal of Health-system Pharmacy* 2001; **58**(18):1693, 1696.
- NCT00149474** *{published data only}*
NCT00149474. Peak flow monitoring in older adults with asthma. <https://clinicaltrials.gov/NCT00149474> (accessed 2 June 2015).
- NCT00232557** *{published data only}*
NCT00232557. Telecommunications system in asthma. www.clinicaltrials.gov/NCT00232557 (accessed 14 August 2015).
- NCT00411346** *{published data only}*
NCT00411346. Patient research In self-management of asthma (PRISMA). <https://clinicaltrials.gov/NCT00411346> (accessed 2 June 2015).
- NCT00562081** *{published data only}*
NCT00562081. The virtual asthma clinic. <https://clinicaltrials.gov/NCT00562081> (accessed 2 June 2015).
- NCT00910585** *{published data only}*
NCT00910585. Coaching in childhood asthma. <https://clinicaltrials.gov/NCT00910585> (accessed 2 June 2015).
- NCT00964301** *{published data only}*
NCT00964301. Telemedicine education for rural children with asthma. <https://clinicaltrials.gov/NCT00964301> (accessed 2 June 2015).
- NCT01117805** *{published data only}*
NCT01117805. Women of color and asthma control. <https://clinicaltrials.gov/NCT01117805> (accessed 2 June 2015).
- Neville 1996** *{published data only}*
Neville RG. Computer assisted assessment and management of patients with asthma. Assessment for a computer assisted assessment and management programme for asthma care. The aim will be to assess if the program will allow clinicians to look at possible future morbidity for each of their patients and with this data discuss and recommend changes in their management of their patient 1996; Vol. 30 September 1998 End date:Publications. Healthcare evaluation.
- Osman 1997** *{published data only}*
* Osman L. A controlled study evaluating relative benefits of two types of review after an A&E attendance. CRISP (Computer Retrieval of Information on Scientific Projects) 1997; Vol. 01 March 2001 End date.
Osman L. A randomised controlled trial of benefits of specialist review or telephone follow up after an Accident & Emergency attendance for asthma. <http://www.isrctn.com/ISRCTN11122844> (accessed 14 August 2015).
- Ostojic 2005** *{published data only}*
* Ostojic V, Cvoricsec B, Ostojic SB, Reznikoff D, Stipic-Markovic A, Tadjman Z. Improving asthma control through telemedicine: A study of short-message service. *Telemedicine Journal and e-Health* 2005;**11**(1):28–35.
Ostojic V, Cvoricsec B, Ostojic SB, Tadjman Z, Stipic-Markovic A. Using of short message service (SMS) for long-term management of asthmatic patients [Abstract]. European Respiratory Society 12th Annual Congress; 2002 Sep 14-18; Stockholm. 2002:Abstract number: P1003.
- Pedram 2012** *{published data only}*
Pedram RS, Piroozmand N, Zolfaghari M, Kazemnejad A, Firoozbakhsh S. Education of how-to-use peak flow meter and following up via SMS on asthma self-management [Farsi]. *Journal of HAYAT* 2012;**18**(4):19–27.
- Peruccio 2005** *{published data only}*
Peruccio D, Bauer B, Delaronde S. Improving asthma treatment and quality of life in a managed care population [Abstract]. *Journal of Allergy and Clinical Immunology* 2005; **115**:S63.
- Petrie 2012** *{published data only}*
Petrie KJ, Perry K, Broadbent E, Weinman J. A text message programme designed to modify patients' illness

and treatment beliefs improves self-reported adherence to asthma preventer medication. *British Journal of Health Psychology* 2012;**17**(1):74–84.

Prabhakaran 2009 {published data only}

* Prabhakaran L, Chee J, Chua K C, Mun WW. The use of text messaging to improve asthma control: a study of short message service (SMS) [Abstract]. *Respirology* 2009; **14**:A217 [PD 10-01].

Prabhakaran L, Chee WY, Chua KC, Abisheganaden J, Wong M. The use of text messaging to improve asthma control: a pilot study using the mobile phone short messaging service (SMS). *Journal of Telemedicine and Telecare* 2010;**16**(5):286–90.

Price 2007 {published data only}

* Price D. A validation of the UK English language Asthma Control Test (ACT) for internet use among adults with asthma. East Norfolk and Waveney Research Consortium (Norfolk & Norwich UH/ Norwich PCT/James Paget/ NWMHP) 2007; Vol. 1 January 2008:Ongoing.

Price D. Validating asthma control test on the internet. Suffolk and Norfolk Research and Development Consortium (S.A.N.D.) 2007; Vol. 30 December 2007: Ongoing.

Raat 2007 {published data only}

Raat H, Mangunkusumo RT, Mohangoo AD, Juniper EF, Van Der Lei J. Internet and written respiratory questionnaires yield equivalent results for adolescents. *Pediatric Pulmonology* 2007;**42**(4):357–61.

Rand 2005 {published data only}

NCT00233233. Assessment mode and validity of self-reports in adults. <https://clinicaltrials.gov/ct2/show/NCT00233233> (accessed 14 August 2015).

Rand CS. Adherence intervention for minority children with asthma. CRISP (Computer Retrieval of Information on Scientific Projects) 2000; Vol. 31 March 2005 End date.

Ricci 2001 {published data only}

Ricci A, Barosi G, Gelmetti A, Esposito R, Tzialla C, Napoli A, et al. A system of teleassistance for in-house monitoring respiratory function in children with bronchial asthma. *Minerva Pediatrica* 2001;**53**(5):466–7.

Rikkens-Mutsaerts 2012 {published data only}

Rikkens-Mutsaerts E, Winters AE, Bakker MJ, van Stel HF, van der Meer V, de Jongste JC, et al. Internet-based self-management compared with usual care in adolescents with asthma: A randomised controlled trial [Abstract]. European Respiratory Society 20th Annual Congress; 2010 Sep 18–22; Barcelona. 2010:[5396].

* Rikkens-Mutsaerts ER, Winters AE, Bakker MJ, van Stel HF, van der Meer V, de Jongste JC, et al. Internet-based self-management compared with usual care in adolescents with asthma: a randomized controlled trial. *Pediatric Pulmonology* 2012;**47**(12):1170–9.

Rosenzweig 2008 {published data only}

Rosenzweig JC, Williams AE, Kite A, Yang M, Kosinski M. Reliability and validity of the asthma control test (ACT) paper and telephone formats [Abstract]. European

Respiratory Society 18th Annual Congress; 2008 Oct 3–7; Berlin. 2008:[P3501].

Ryan 2012 {published data only}

Ryan D, Pinnock H, Amanda L, Lionel T, Sheikh A, Musgrave S, et al. Can mobile phone technology improve asthma control? A randomised trial [Abstract]. European Respiratory Society 20th Annual Congress; 2010 Sep 18–22; Barcelona. 2010:[2138].

Ryan D, Pinnock H, Lee AJ, Tarassenko L, Ayansina D, Musgrave S, et al. Can your mobile phone help your asthma: preliminary results [Abstract]. *Primary Care Respiratory Journal* 2010;**19**:A15 [54].

Ryan D, Pinnock H, Lee AJ, Tarassenko L, Pagliari C, Sheikh A, et al. The CYMPLA trial. Mobile phone-based structured intervention to achieve asthma control in patients with uncontrolled persistent asthma: A pragmatic randomised controlled trial. *Primary Care Respiratory Journal* 2009;**18**(4):343–5.

Ryan D, Pinnock H, Tarassenko L, Lee A, Sheikh A, Price D. Can your mobile phone improve your asthma? [Abstract]. *Thorax* 2010;**65**:S135.

* Ryan D, Price D, Musgrave SD, Malhotra S, Lee AJ, Ayansina D, et al. Clinical and cost effectiveness of mobile phone supported self monitoring of asthma: multicentre randomised controlled trial. *BMJ* 2012;**344**:e1756.

Schatz 2003 {published data only}

Schatz M, Rodriguez E, Falkoff R, Zeiger RS. The relationship of frequency of follow-up visits to asthma outcomes in patients with moderate persistent asthma. *Journal of Asthma* 2003;**40**(1):49–53.

Schatz 2010 {published data only}

Schatz M, Zeiger RS. Ineffectiveness of telephone-based environmental control intervention to improve asthma outcomes. *Journal of Allergy and Clinical Immunology* 2010; **126**(4):873–5.

Searing 2012 {published data only}

Searing DA, Bender BG. Short message service (SMS) for asthma management: A pilot study utilizing text messaging to promote asthma self-management [Abstract]. Annual Meeting of American Academy of Allergy, Asthma and Immunology (AAAAI); 2012 March 2–6; Orlando. 2012; Vol. 129:AB142 [538].

Seid 2012 {published data only}

Seid M, D'Amico EJ, Varni JW, Munafò JK, Britto MT, Kercksmar CM, et al. The in vivo adherence intervention for at risk adolescents with asthma: report of a randomized pilot study. *Journal of Pediatric Psychology* 2012;**37**(4):390–403.

Shanovich 2009 {published data only}

Shanovich KK, Pulvermacher AD, Hollman SJ, Richardson PA, Wise ME, Lee SH, et al. Nurse case management services provided to supplement a web-based asthma education program [Abstract]. *Journal of Allergy and Clinical Immunology* 2008;**121**:S40 [155].

* Shanovich KK, Sorkness CA, Wise ME, Pulvermacher AD, Bhattacharya A, Gustafson DH. Internet Telehealth for Pediatric Nurse Case Management Improves Asthma

- Control [Abstract]. *Journal of Allergy and Clinical Immunology* 2009;**123**:S43.
- Taitel 2014** *{published data only}*
Taitel M, Jiang JZ, Rudkin K. Impact of pharmacist-call intervention program on new-to-therapy patients' medication adherence. *Value in Health* 2014;**17**:A138–9.
- Uysal 2013** *{published data only}*
Uysal MA, Mungan D, Yorgancioglu A, Yildiz F, Akgun M, Gemicioglu B, et al. Asthma control test via text messaging: could it be a tool for evaluating asthma control?. *Journal of Asthma* 2013;**50**(10):1083–9.
- van den Berg 2002** *{published data only}*
van den Berg NJ, ten Have WH, Bindels PJE, van der Palen J, van Aalderen WMC. Is the availability of a 24 hour asthma-telephone useful in the implementation of asthma treatment guidelines for children aged 6-16 years, among general practitioners? [Abstract]. *European Respiratory Journal* 2002;**20**:329s.
- van der Meer 2009** *{published data only}*
van der Meer V, Bakker MJ, Rabe KF, Sterk PJ, Kievit J, Assendelft WJJ, et al. Improved quality of life by internet based self management versus usual care in asthma a randomized controlled trial [Abstract]. European Respiratory Society 18th Annual Congress; 2008 Oct 3-7; Berlin. 2008:[1372].
van der Meer V, Bakker MJ, van den Hout WB, Rabe KF, Sterk J, Kievit J, et al. Self-management for asthma on the Internet: a randomized study. *Nederlands Tijdschrift Voor Geneeskunde* 2010;**154**:403–9.
* van der Meer V, Bakker MJ, van den Hout WB, Rabe KF, Sterk PJ, Kievit J, et al. Internet-based self-management plus education compared with usual care in asthma: A randomized trial. *Annals of Internal Medicine* 2009;**151**(2): 110–20.
van der Meer V, van Stel HF, Bakker MJ, Roldaan AC, Assendelft WJ, Sterk PJ, et al. Weekly self-monitoring and treatment adjustment benefit patients with partly controlled and uncontrolled asthma: an analysis of the SMASHING study. *Respiratory Research* 2010;**11**:74.
van der Meer V, van den Hout WB, Bakker MJ, Rabe KF, Sterk PJ, Assendelft WJJ, et al. Cost-effectiveness of internet-based self-management compared with usual care in asthma. *PLoS One* 2011;**6**(11):e27108.
van der Meer, van Stel HF, Bakker MJ, Roldaan AC, Assendelft WJJ, Sterk PJ, et al. Internet-based self management improves short-term asthma control: the SMASHING study [Abstract]. *Primary Care Respiratory Journal* 2008;**17**:124.
- van Gaalen 2012** *{published data only}*
* van Gaalen JL, Bakker MJ, van Bodegom-Vos L, Snoeck-Stroband JB, Assendelft WJ, Kaptein AA, et al. Implementation strategies of internet-based asthma self-management support in usual care. Study protocol for the IMPASSE cluster randomized trial. *Implementation Science* 2012;**7**:113.
van Gaalen JL, Beerthuizen T, van der Meer V, van Reisen P, Redelijkheid GW, Snoeck-Stroband JB, et al. Long-term outcomes of internet-based self-management support in adults with asthma: randomized controlled trial. *Journal of Medical Internet Research* 2013;**15**(9):e188.
- van Reisen 2010** *{published data only}*
van Reisen PAJ, Redelijkheid WG, van Gaalen JL, van der Meer V, Bakker MJ, Beerthuizen T, et al. The long-term outcomes of a one-year Internet-based self-management support programme compared with usual care in asthma. Additional follow-up 1.5 year after a randomized trial [Abstract]. European Respiratory Society 20th Annual Congress; 2010 Sep 18-22; Barcelona. 2010:[E2160].
- Vasbinder 2013** *{published data only}*
Vasbinder EC, Janssens HM, Rutten-van Mólken MP, van Dijk L, de Winter BC, de Groot RC, et al. e-Monitoring of Asthma Therapy to Improve Compliance in children using a real-time medication monitoring system (RTMM): the e-MATIC study protocol. *BMC Medical Informatics and Decision Making* 2013;**13**:38.
- Vollmer 2006** *{published data only}*
* Vollmer WM, Kirshner M, Peters D, Drane A, Stibolt T, Hickey T, et al. Use and impact of an automated telephone outreach system for asthma in a managed care setting. *American Journal of Managed Care* 2006;**12**(12):725–33.
Vollmer WM, Peters D, Kirshner M, Naleway A, Buist AS, Hickey T, et al. Use of a telephone triage system for asthma management [Abstract]. American Thoracic Society 2005 International Conference; May 20-25; San Diego. 2005: [A40] [Poster: G15].
- Voorend-van Bergen 2013** *{published data only}*
Voorend-van Bergen S, Vaessen-Verberne A, Landstra A, Brackel H, van den Berg N, de Jongste J, et al. FeNO and web-based monitoring in paediatric asthma management; the BATMAN study [Abstract]. *European Respiratory Journal* 2013;**42**(Suppl 57):629s [3014].
- Wiecha 2007** *{published data only}*
Wiecha JM, Adams WG. BostonBreathes: an RCT to improve pediatric asthma care with a home-based interactive website for patient education, monitoring, and clinical teamwork. AMIA Annual Symposium Proceedings. 2007: 1154.
- Willems 2008** *{published data only}*
* Willems DC, Joore MA, Hendriks JJ, Nieman FH, Severens JL, Wouters EF. The effectiveness of nurse-led telemonitoring of asthma: results of a randomized controlled trial. *Journal of Evaluation in Clinical Practice* 2008;**14**(4):600–9.
Willems DC, Joore MA, Hendriks JJ, van Duurling RA, Wouters EF, Severens JL. Process evaluation of a nurse-led telemonitoring programme for patients with asthma. *Journal of Telemedicine & Telecare* 2007;**13**(6):310–7.
Willems DCM, Joore MA, Hendriks JJE, Wouters EFM, Severens JL. Cost-effectiveness of a nurse-led telemonitoring intervention based on peak expiratory flow measurements in asthmatics: results of a randomised controlled trial. *Cost Effectiveness and Resource Allocation* 2007;**5**:10.

Young 2012 *[published data only]*

Young HN, Havican SN, Griesbach S, Thorpe JM, Chewning BA, Sorkness CA. Patient and pharmacist telephonic encounters (PARTE) in an underserved rural patient population with asthma: results of a pilot study. *Telemedicine Journal and e-Health* 2012;**18**(6):427–33.

Yun 2013 *[published data only]*

Yun TJ, Arriaga RI. A text message a day keeps the pulmonologist away. Proceedings of the SIGCHI conference on human factors in computing systems; 2013 April 27-May 02; Paris. 2013:1769–78.

Zachgo 2002 *[published data only]*

Zachgo W, Juenemann R, Halbich C, Hofmann N, Eleljanova S, Reiss Plagemann C, et al. Optimized computer-based inhaled steroid therapy in asthma. *American Journal of Respiratory and Critical Care Medicine* 2002;**165**:A766.

Additional references**ATS 2000**

American Thoracic Society. Proceedings of the ATS workshop on refractory asthma: current understanding, recommendations, and unanswered questions. *American Journal of Respiratory and Critical Care Medicine* 2000;**162**(6):2341–51.

BTS/SIGN 2014

British Thoracic Society/Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma, October 2014. <https://www.brit-thoracic.org.uk/guidelines-and-quality-standards/asthma-guideline/> (accessed 27 February 2015).

Department of Health 2012

Department of Health. Long term conditions compendium of information: third edition (30 May 2012). <https://www.gov.uk/government/publications/long-term-conditions-compendium-of-information-third-edition> (accessed 3 March 2015).

Department of Health 2013

Department of Health. A mandate from the Government to the NHS Commissioning Board: April 2013 to March 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/256497/13-15_mandate.pdf (accessed 27 February 2015).

Eysenback 2001

Eysenback G. What is e-health?. *Journal of Medical Internet Research* 2001;**3**(2):e20.

Gibson 2002

Gibson PG, Powell H, Wilson A, Abramson MJ, Haywood P, Bauman A, et al. Self-management education and regular practitioner review for adults with asthma. *Cochrane Database of Systematic Reviews* 2002, Issue 3. [DOI: 10.1002/14651858.CD001117]

GINA 2014

Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. www.ginaasthma.org/local/

[uploads/files/GINA_Report_2014_Aug12.pdf](https://www.ginaasthma.org/local/uploads/files/GINA_Report_2014_Aug12.pdf) (accessed 27 February 2015).

Global Asthma Report 2014

International Union Against Tuberculosis and Lung Disease. The Global Asthma Report 2014. <http://theunion.org/what-we-do/publications/technical/global-asthma-report> (accessed 30 April 2015).

GRADEpro GDT 2015 [Computer program]

McMaster University (developed by Evidence Prime Inc.). GRADEpro Guideline Development Tool. Available from www.gradepr.org. McMaster University (developed by Evidence Prime Inc.), 2015.

Higgins 2011

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Jaana 2009

Jaana M, Paré G, Sicotte C. Home telemonitoring for respiratory conditions: a systematic review. *American Journal of Managed Care* 2009;**15**(5):313–20.

Jannett 2003

Jennett PA, Affleck Hall L, Hailey D, Ohinmaa A, Anderson C, Thomas R, et al. The socio-economic impact of telehealth: a systematic review. *Journal of Telemedicine and Telecare* 2003;**9**(6):311–20.

Juniper 1994

Juniper EF, Guyatt GH, Willan A, Griffith LE. Determining a minimal important change in a disease-specific quality of life questionnaire. *Journal of Clinical Epidemiology* 1994;**47**(1):81–7.

Juniper 1999

Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *European Respiratory Journal* 1999;**14**(4):902–7.

Kew 2015a

Kew K, Cates C. Asthma monitoring with remote feedback from a health professional. *Cochrane Database of Systematic Reviews* 2015, Issue 5. [DOI: 10.1002/14651858.CD011714]

Laver 2013

Laver KE, Schoene D, Crotty M, George S, Lannin NA, Sherrington C. Telerehabilitation services for stroke. *Cochrane Database of Systematic Reviews* 2013, Issue 12. [DOI: 10.1002/14651858.CD010255.pub2]

Lustig 2012

Lustig TA, Board on Health Care Services, Institute of Medicine. *The Role of Telehealth in an Evolving Health Care Environment: Workshop Summary*. Washington, DC: The National Academies Press, 2012.

McLean 2010

McLean S, Chandler D, Nurmatov U, Liu JLY, Pagliari C, Car J, et al. Telehealthcare for asthma. *Cochrane Database*

of *Systematic Reviews* 2010, Issue 10. [DOI: 10.1002/14651858.CD007717.pub2]

McLean 2011

McLean S, Nurmatov U, Liu JLY, Pagliari C, Car J, Sheikh A. Telehealthcare for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2011, Issue 7. [DOI: 10.1002/14651858.CD007718.pub2]

McLean 2013

McLean S, Sheikh A, Cresswell K, Nurmatov U, Mukherjee M, Hemmi A, et al. The impact of telehealthcare on the quality and safety of care: a systematic overview. *PLoS One* 2013;**8**(8):e71238.

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman D, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* 2009;**6**(7):e1000097. [DOI: 10.1371/journal.pmed.1000097]

NRAD 2014

National Review of Asthma Deaths. Why asthma still kills: the national review of asthma deaths. <https://www.rcplondon.ac.uk/sites/default/files/why-asthma-still-kills-full-report.pdf> (accessed 27 February 2015).

Pinnock 2007b

Pinnock H, Slack R, Pagliari C, Price D, Sheikh A. Understanding the potential role of mobile phone-based

monitoring on asthma self-management: qualitative study. *Clinical and Experimental Allergy* 2007;**37**(5):794–802.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Steventon 2012

Steventon A, Bardsley M, Billings J, Dixon J, Doll H, Hirani S, et al. Effect of telehealth on use of secondary care and mortality: findings from the Whole System Demonstrator cluster randomised trial. *BMJ* 2012;**344**:e3874.

Zhao 2014

Zhao J, Zhai YK, Zhu WJ, Sun DX. Effectiveness of telemedicine for controlling asthma symptoms: a systematic review and meta-analysis. *Telemedicine Journal and e-Health* 2014;**21**(6):484–92.

References to other published versions of this review

Kew 2015b

Kew KM, Cates CJ. Remote versus face-to-face asthma reviews. *Cochrane Database of Systematic Reviews* 2015, Issue 5. [DOI: 10.1002/14651858.CD011715]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Chan 2007

Methods	<p>Study design: 12 month parallel RCT</p> <p>Setting: paediatric clinic at Tripler Army Medical Center, Hawaii</p> <p>Enrolment began in March 2003 and ended in December 2003. Participant data collection ended with the last participant's final visit in February 2005</p>	
Participants	<p>Population: 120 children were randomised to the virtual group (60) or the office-based group (60)</p> <p>Baseline characteristics:</p> <p>mean age, years (SD): remote 10.2 (3.1); face-to-face 9.0 (3.0)</p> <p>% male: remote 61.7; face-to-face 63.3</p> <p>% predicted FEV₁ (SD): remote 104.1 (19.9); face-to-face 96.8 (13.0)</p> <p>Inclusion criteria: children aged 6 to 17 years with persistent asthma, dependent of active duty or retired USA military personnel, not moving from Oahu for 12 months after entry into the study, ability to receive cable modem connections in the home, willingness to complete questionnaires and monitoring</p> <p>Exclusion criteria: none stated</p>	
Interventions	<p>Intervention: virtual group participants received computers, internet connections and in-home internet-based case management and received education through the study website</p> <p>Control: office-based group patients received traditional in-person education and case management</p>	
Outcomes	<p>Control medication use, daily symptom diary, peak flow, patient and caregiver AQLQ, service utilisation, asthma knowledge retention</p> <p>Measured at 2 weeks, 6 weeks, 3 months, 6 months and 12 months</p>	
Notes	<p>Funding: grant from the US Army Medical Research Acquisition Activity</p> <p>ID number(s): N/A</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"patients underwent block randomisation with a table of random numbers"
Allocation concealment (selection bias)	Unclear risk	The study did not provide any details.
Blinding of participants and personnel Objective outcomes	Low risk	It would not have been possible to blind participants and personnel to allocation due to the nature of the intervention. However, participants and personnel being aware of group allocation is unlikely to have

Chan 2007 (Continued)

		affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (Asthma Control Questionnaire (ACQ) and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	It is possible to blind outcome assessment but the study did not provide any specific details of whether this was done
Incomplete outcome data (attrition bias) All outcomes	High risk	Dropout was much higher in the virtual group (23%) than the office group (8%). The study authors did not account for non-adherent participants and other dropouts in the analysis
Selective reporting (reporting bias)	Low risk	Outcomes were well reported. There was no protocol registration available to check all pre-specified measures were included but there was no evidence of selective reporting
Other bias	Low risk	We did not note any other possible sources of bias.

Gruffydd-Jones 2005

Methods	Study design: 12 month parallel RCT Setting: 1 practice in England, UK Participants were recruited between December 2002 and March 2003
Participants	Population: 194 people were randomised to the telephone group (97) or the clinic group (97) Baseline characteristics: mean age, years (standard deviation (SD)): remote 50.8 (15.4); face-to-face 49.6 (16.1) % male: remote 51.5; face-to-face 39.2 % predicted forced expiratory volume in one second (FEV ₁) (SD): not reported Inclusion criteria: adults with asthma aged 17 to 70 years and on the practice asthma list Exclusion criteria: housebound, did not possess a telephone or were unwilling to give informed consent
Interventions	Intervention: participants were contacted by telephone at 6-monthly intervals by 1 of 2 trained asthma nurses. The participant was then asked the RCPs 'three questions' plus two extra questions related to a high risk of asthma death. The nurse formulated an

	individualised asthma action plan with the participant, with advice on what to do if asthma control deteriorated Control: participants received usual care by 6-monthly check up via a dedicated asthma appointment with a diploma-level asthma nurse. Symptom scores, inhaler technique and peak flow measurements were checked and all participants issued with an asthma action plan	
Outcomes	ACQ, mini-AQLQ, mild and severe exacerbations, healthcare costs, clinical time, inhaler use, unscheduled healthcare visits all given per patient year Measured at baseline, 6 months and 12 months	
Notes	Funding: grant from Asthma UK ID number(s): N/A	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The study randomised participants using a random number tables on a 1 to 1 basis and stratified according to severity
Allocation concealment (selection bias)	Unclear risk	The study did not provide any details.
Blinding of participants and personnel Objective outcomes	Low risk	It would not have been possible to blind participants and personnel to allocation due to the nature of the intervention. However, participants and personnel being aware of group allocation is unlikely to have affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (ACQ and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	High risk	"assessors were not blinded to the interventions due to limited resources"
Incomplete outcome data (attrition bias) All outcomes	High risk	"There were 20 withdrawals in the control group after the first visit, mainly due to non-attendance and 6 in the telephone group, one of which was due to non-attendance. As this trial is as real-world as possible the fact that there was a high non-attendance rate was taken account of in

		analysing the costs.“
Selective reporting (reporting bias)	Low risk	Outcomes were well reported. There was no protocol registration available to check all pre-specified measures were included but there was no evidence of selective reporting
Other bias	Low risk	We did not note any other possible sources of bias.

Hashimoto 2011

Methods	<p>Study design: pragmatic 6 month parallel RCT</p> <p>Setting: 2 academic tertiary care hospitals and 4 large community hospitals in The Netherlands</p> <p>Participants were randomised between November 2007 and October 2008</p>
Participants	<p>Population: 95 people were randomised to the internet group (52) or the conventional face-to-face management group (38)</p> <p>Baseline characteristics:</p> <p>mean age, years (SD): remote 48.5 (12.4); face-to-face 52.4 (11.7)</p> <p>% male: remote 45; face-to-face 47</p> <p>% predicted FEV₁ (SD): remote 76.3 (24.7); face-to-face 71.3 (21.0)</p> <p>Inclusion criteria: adults (18 to 75 years) with a diagnosis of severe refractory asthma according to the major and minor criteria recommended by the American Thoracic Society. They had uncontrolled asthma despite intensive follow-up by an asthma specialist for at least 1 year, chronic treatment with oral corticosteroids and high doses of ICS plus long-acting bronchodilators. All were non-smokers with a maximum smoking history of 15 pack-years and had access to internet or mobile telephone</p> <p>Exclusion criteria: none stated</p>
Interventions	<p>Intervention: dose adjustment of oral corticosteroids guided by an internet-based management tool (internet group). Included electronic diary, decision support and monitoring support by a study nurse</p> <p>Control: dose adjustment of oral corticosteroids according to conventional asthma treatment by the pulmonologist, according to GINA (conventional management group)</p>
Outcomes	<p>Cumulative sparing of oral corticosteroid therapy (OCS), ACQ, AQLQ, global satisfaction scale, FEV₁, number of exacerbations and days of hospitalisation</p> <p>The authors defined an exacerbation as a decrease in morning FEV₁ of at least 10% compared with the mean FEV₁ from the week before, or a respiratory event requiring an increase in prednisone equivalent to at least 10 mg/day, or a course of antibiotics, with or without hospitalisation</p>
Notes	<p>Funding: The Netherlands Organization for Health Research and Development (ZonMw)</p> <p>ID number(s): 1146 (Netherlands Trial Reg No.)</p>

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The study randomised participants by a computer random number generator and remained on the same allocation throughout the study. Communication: "The random codes were stratified for study center and initial dose of corticosteroid dose (lower or higher than 10 mg prednisone per day)"
Allocation concealment (selection bias)	Unclear risk	"unblinded after randomisation", implies it was concealed, but the study did not provide any further details
Blinding of participants and personnel Objective outcomes	Low risk	The treatment assignments were unblinded after randomisation to allow monthly corticosteroid dose adjustments according to conventional treatment by the physician or weekly adjustments according to the internet algorithm. While it was not possible to blind participants and personnel to allocation due to the nature of the intervention, participants and personnel being aware of group allocation is unlikely to have affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (ACQ and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Communication: "This was a pragmatic study so the outcome assessors were not blind to the group allocation in order to allow monthly corticosteroid dose adjustments (according to conventional treatment by the physician) or weekly adjustments (according to the internet algorithm)"
Incomplete outcome data (attrition bias) All outcomes	High risk	Five participants in the conventional management group withdrew consent before the study had started and the study ex-

Hashimoto 2011 (Continued)

		cluded one participant because of poor adherence to the trial protocol. The study included 89 participants out of 95 in the intention-to-treat (ITT) analysis; 51 and 38. Dropout was higher in the conventional treatment group (16% versus 8%)
Selective reporting (reporting bias)	Low risk	The study was prospectively registered, and outcomes were well reported
Other bias	Low risk	We did not note any other possible sources of bias.

Pinnock 2003

Methods	<p>Study design: pragmatic parallel RCT (duration of study participation varied across participants)</p> <p>Setting: 4 general practices in the UK</p>
Participants	<p>Population: 278 people were randomised to remote telephone check-up (137) or face-to-face check-up (141)</p> <p>Baseline characteristics: mean age, years (SD): remote 54.6 (17.5); face-to-face 56.4 (17.5) % male: remote 41; face-to-face 42 % predicted FEV₁ (SD): not reported</p> <p>Inclusion criteria: adults with asthma who had requested a prescription for a bronchodilator inhaler in the last 6 months</p> <p>Exclusion criteria: if diagnosis of asthma had been made within the previous year, if they had chronic obstructive pulmonary disease, if communication difficulties made a telephone check-up impossible, or (at the general practitioner's (GP's) request) for major social or medical reasons</p>
Interventions	<p>Intervention: telephone check-up with the asthma nurse. The nurse tried up to 4 times to contact the participants</p> <p>Control: face-to-face check-ups in the surgery also with the asthma nurse, one invitation was sent in the usual manner. Content of the check-up was as the nurse deemed appropriate</p>
Outcomes	Medical reviews, time taken to review participants in each arm, asthma morbidity on the short Q, asthma related quality of life on the mini AQLQ, participant satisfaction, costs
Notes	<p>Funding: originally developed at a General Practice Airways Group research meeting, which was organised by Mark Levy and funded by an educational grant from AstraZeneca. The trial was funded by British Lung Foundation (Grant No P00/9). Additionally, one study author was supported by an NHS R&D national primary care fellowship.</p> <p>ID number(s): N/A</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were centrally randomised in blocks of 10 to ensure that approximately equal numbers of participants were allocated to each study arm
Allocation concealment (selection bias)	Low risk	"Centrally randomised" implies that allocation was undertaken independently and concealed
Blinding of participants and personnel Objective outcomes	Low risk	It would not have been possible to blind participants and personnel to allocation due to the nature of the intervention. However, participants and personnel being aware of group allocation is unlikely to have affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (ACQ and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	"a researcher, blinded to allocation visited each of the practices and validated a random 20% sample of consultation data and data retrieved from records". However, the participants and investigators could not be blinded to the interventions and, in most cases, the outcome assessors were not blinded to group allocation either
Incomplete outcome data (attrition bias) All outcomes	Low risk	The percentage of withdrawals was low and even between groups (5.1 and 4.3% in the remote and face-to-face groups respectively)
Selective reporting (reporting bias)	Low risk	There was no evidence of selective reporting.
Other bias	Low risk	We did not note any other possible sources of bias.

Pinnock 2007a

Methods	<p>Study design: 12 month before-and-after implementation study</p> <p>Setting: 1 large English general practice on 3 sites</p>
Participants	<p>Population: 3 practices were randomised to: 1. a choice of remote phone check-ups or face-to-face check-ups (554 on list), 2. face-to-face only check-ups (659 on list), or 3. a usual care control group which was not included in this systematic review (515 on list)</p> <p>Baseline characteristics: mean age, years (SD): remote 43 (24.8); face-to-face 42.3 (24.4) % male: remote 44.2; face-to-face 44.9 % predicted FEV₁ (SD): not reported</p> <p>Inclusion criteria: adults and adolescents with a diagnosis of asthma and prescribed asthma medication in the previous year</p> <p>Exclusion criteria: children under 12 years of age, diagnosis of COPD</p>
Interventions	<p>Intervention: participants were identified from the practice computer database and sent 3 invitations over the study period. They could book either a telephone or face-to-face check-up both at a pre-arranged time. Participants who did not respond to the 3 invitations were phoned and reviewed opportunistically</p> <p>Control: participants were recalled to face-to-face only asthma check-ups using invitations by post or as memos with repeat prescriptions. There was no option of telephone check-ups and no systematic attempt was made to phone non-attenders opportunistically</p> <p>Group excluded: the usual-care control group maintained their well established asthma clinic, and existing procedures (for example, invitations are issued in response to clinical need), but no systematic recall was undertaken</p>
Outcomes	Proportion reviewed, asthma morbidity and enablement on the mini AQLQ, ACQ, modified patient enablement instrument and Asthma Bother Profile, adverse events, costs
Notes	<p>Funding: Scientific Foundation Board of the Royal College of General Practitioners (SFB/2003/45)</p> <p>ID number(s): N/A</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The study decided allocation to the telephone option by the public toss of a coin
Allocation concealment (selection bias)	High risk	The study allocated individuals to treatment after the two clusters had been decided by the toss of a coin
Blinding of participants and personnel Objective outcomes	Low risk	It would not have been possible to blind participants and personnel to allocation due to the nature of the intervention. However, participants and personnel being

Pinnock 2007a (Continued)

		aware of group allocation is unlikely to have affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (ACQ and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	The nurses were aware of allocation but it was unclear whether it was the nurses measuring outcomes, or if it was someone independent from the study who could remain blind to allocation. The study did not describe this. The study stated that there were quality control checks blinded to allocation which confirmed accuracy of data transfer
Incomplete outcome data (attrition bias) All outcomes	Low risk	Real-world implementation study, therefore the uptake rate by participants is part of the study, routine asthma check-up was provided for 66.3% of participants in the telephone only group and 53.8% in the face-to-face only group
Selective reporting (reporting bias)	Low risk	There was no evidence of selective reporting.
Other bias	High risk	This study was a cluster implementation study with a before-and-after design. It randomised 2 practices to the interventions which would not have controlled for baseline imbalances in the same way as individual randomisation, and this meant the participant population in each group was not static. The intervention was a telephone option and many in that practice opted for a usual face-to-face check-up, which meant the study was not making a direct comparison of remote and face-to-face check-ups. Additionally, people in the telephone option group were phoned opportunistically to increase uptake of check-ups which did not happen in the face-to-face group. These factors mean we cannot be certain that mode of check-up, and not the increased likelihood of check-up, was the variable being measured

Rasmussen 2005

Methods	Study design: 6 month pragmatic parallel RCT Setting: general practices and specialist clinics in Copenhagen, Denmark
Participants	Population: 300 people were randomised to remote check-ups (100), face-to-face check-ups with a specialist (100), and a usual care group not included in this review (100) Baseline characteristics: mean age, years (SD): remote 28 (NR); face-to-face 30 (NR) % male: remote 31.8; face-to-face 34.1 % predicted FEV ₁ (SD): remote 91 (NR); face-to-face 93 (NR) Inclusion criteria: 18 to 45 years with definite asthma, living in the catchment area of H:S Bispebjerg University Hospital of Copenhagen, Denmark Exclusion criteria: none stated
Interventions	Intervention: participants were given a Peak Flow Meter and taught how to fill in a daily diary and respond to the computer's advice. Physicians gave instructions via e-mail or telephone to the participant. The intervention included an electronic diary, an asthma action plan and a decision support system for the physician Control: the specialists taught the participants how to adjust their medication on the basis of a peak flow meter and written action plan Group not included: the GP group was asked to contact their GP and pass on a letter describing the study and giving the test results. GPs in Copenhagen had been sent a circular about asthma and GINA guidelines in the past
Outcomes	AQLQ, asthma self-care, smoking, education, salary, sick leave, hospitalisations, medication compliance, adverse events, lung function Measured at baseline and 6 months
Notes	Funding: Grants from H:S Corporation of University Hospital of Copenhagen, AstraZeneca, and private funds ID number(s): N/A

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Communication: "The allocation sequences were computer-generated by a senior respiratory physician. These sequences consisted of randomised blocks of 30 asthmatics"
Allocation concealment (selection bias)	Low risk	Communication: "The envelopes were packed by two medical students one month before the start of the study and the randomisation lists were stored in a separate, sealed envelope. The consecutively numbered and sealed envelopes contained the randomisation code. All envelopes were opened sequentially after the asthma diagnosis had been verified"

Blinding of participants and personnel Objective outcomes	Low risk	It would not have been possible to blind participants and personnel to allocation due to the nature of the intervention. However, participants and personnel being aware of group allocation is unlikely to have affected the results for the objective outcomes (exacerbations and adverse events)
Blinding of participants and personnel Subjective outcomes	High risk	Participants and personnel being aware of group allocation could have affected their scores on subjective outcomes such as those measured on self-report scales (ACQ and AQLQ)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Communication: "It was not possible to blind outcome assessors to group allocation"
Incomplete outcome data (attrition bias) All outcomes	High risk	Of the 300 participants randomised, 253 participants completed both the screening and follow-up visits. Dropout was unbalanced across groups (12%, 15% and 20%) , and the study does not appear to have imputed data for missing values
Selective reporting (reporting bias)	Low risk	The paper did not report all of the results from the questionnaires but the lead study author provided them on request
Other bias	Low risk	We did not note any other possible sources of bias.

Abbreviations: ACQ = Asthma Control Questionnaire; AQLQ = Asthma Quality of Life Questionnaire; COPD = chronic obstructive pulmonary disease; FEV₁ = forced expiratory volume in one second; GINA = Global Initiative for Asthma; GP = general practitioner; ICS = inhaled corticosteroids; ITT = intention-to-treat analysis; NR = not reported; OCS = oral corticosteroids; RCP = respiratory care practitioner; RCT = randomised controlled trial; SD = standard deviation.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
ACTRN12606000400561	Wrong intervention - technology-based self management between reviews
Ahmed 2011	Wrong intervention - technology-based self management between reviews
Andersen 2007	Wrong intervention - minimal or no provider involvement
Apter 2000	Wrong design - not a trial report
Apter 2015	Wrong comparison - telemedicine portal used with or without home visits (both groups used the portal)
Araujo 2012	Wrong design - crossover RCT
Baptist 2013	Wrong comparison - phone calls for asthma education versus non-asthma phone calls
Barbanel 2003	Wrong intervention - asthma education intervention led by a pharmacist
Bateman 2000	Wrong intervention - technology-based self management between reviews
Bender 2001	Wrong intervention - study assessing validity of self-reports
Bender 2007	Wrong intervention - study assessing validity of self-reports
Bender 2010	Wrong intervention - minimal or no provider involvement
Boyd 2014	Wrong intervention - pharmacist led intervention about adherence
Burbank 2012	Wrong intervention - focus on asthma education, not monitoring with remote reviews
Burkhart 2002	Wrong intervention - intervention to improve adherence to home PEF measurements
Bynum 2001	Wrong intervention - pharmacy led technology intervention to improve adherence
Chandler 1990	Wrong intervention - monitoring theophylline levels
Chatkin 2006	Wrong intervention - phone calls to promote adherence, not remote reviews
Chen 2013	Wrong intervention - asthma behavioural intervention using technology, not remote reviews
Cicutto 2009	Wrong intervention - not remote reviews
Clark 2007	Wrong intervention - counselling intervention not remote reviews
Clarke 2014	Wrong intervention - parenting intervention, not remote reviews
Claus 2004	Wrong design - not a randomised controlled trial (RCT)

(Continued)

Cruz-Correia 2007	Wrong design - crossover RCT
de Jongste 2008	Wrong comparison - comparing 2 types of electronic monitoring (FeNo versus symptoms)
De Vera 2014	Wrong intervention - asthma education and adherence monitoring by a pharmacist
Deschildre 2012	Wrong intervention - technology-based self management between reviews
Donald 2008	Wrong intervention - technology-based self management between reviews
Dwinger 2013	Wrong intervention - coaching/education intervention using technology for multiple chronic conditions
Eakin 2012	Wrong intervention - not remote asthma reviews
Finkelstein 2005	Wrong intervention - technology-based self management between reviews
Fonseca 2006	Wrong design - survey of RCT participants
Foster 2014	Wrong intervention - adherence intervention
Friedman 1999	Wrong intervention - mostly automated home monitoring, not remote reviews
Garbutt 2010	Wrong intervention - asthma coaching/education intervention over the phone, not remote reviews
Guendelman 2002	Wrong intervention - technology-based self management between reviews
Gustafson 2012	Wrong intervention - self-determination theory intervention, not remote reviews
Halterman 2012	Wrong intervention - technology-based self management between reviews
Huang 2013	Wrong intervention - support intervention, not remote reviews
Jan 2007	Wrong intervention - technology-based self management between reviews
Janevic 2012	Wrong intervention - management intervention for African American women, not remote reviews
Jerant 2003	Wrong intervention - mixed diagnosis study comparing models of delivering home care
Kattan 2006	Wrong intervention - minimal or no provider involvement
Khan 2003	Wrong intervention - one phone call at discharge, not remote reviews
Kojima 2005	Wrong intervention - not technology-based
Kokubu 1999	Wrong intervention - technology-based self management between reviews

(Continued)

Lam 2011	Wrong design - cross-sectional analysis of an ongoing RCT, and mixed diagnosis
Liu 2011	Wrong intervention - technology-based self management between reviews
Lobach 2013	Wrong intervention - not about remote reviews
McCowan 2001	Wrong intervention - computer-aided decision support during consultation
McPherson 2006	Wrong intervention - asthma education delivered via CD-ROM and book versus book alone
Merchant 2013	Wrong intervention - remote monitoring of inhaler adherence
Morrison 2014	Wrong intervention - minimal or no provider involvement
Murphy 2001	Wrong design - comment on a RCT
NCT00149474	Wrong comparison - remote monitoring using PEF or symptoms
NCT00232557	Wrong comparison - phone monitoring plus asthma education versus phone education
NCT00411346	Wrong intervention - technology-based self management between reviews
NCT00562081	Wrong intervention - focus on asthma education not remote reviews
NCT00910585	Wrong intervention - focus on asthma education not remote reviews
NCT00964301	Wrong intervention - focus on asthma education not remote reviews
NCT01117805	Wrong intervention - counselling not remote reviews
Neville 1996	Wrong intervention - computer-aided decision support during consultation
Osman 1997	Wrong intervention - post admission follow-up
Ostojic 2005	Wrong intervention - technology-based self management between reviews
Pedram 2012	Wrong intervention - main focus of the study was to educate participants on how to use a peak flow meter
Peruccio 2005	Wrong intervention - treatment awareness education delivered over the phone
Petrie 2012	Wrong intervention - minimal or no provider involvement
Prabhakaran 2009	Wrong intervention - technology-based self management between reviews
Price 2007	Wrong intervention - validating the Asthma Control Test for internet use

(Continued)

Raat 2007	Wrong design - questionnaire not a RCT
Rand 2005	Wrong intervention - study measuring validity of self-report
Ricci 2001	Wrong intervention - technology-based self management between reviews
Rijkers-Mutsaerts 2012	Wrong intervention - minimal or no provider involvement
Rosenzweig 2008	Wrong intervention - validation study
Ryan 2012	Wrong intervention - technology-based self management between reviews
Schatz 2003	Wrong comparison - phone calls on top of face-to-face review, not instead of
Schatz 2010	Wrong intervention - letter regarding validation of telephone delivery of the Asthma Control Questionnaire (ACQ)
Searing 2012	Wrong intervention - minimal or no provider involvement
Seid 2012	Wrong intervention - asthma education and motivational interviewing, not remote reviews
Shanovich 2009	Wrong intervention - focus on asthma education not remote reviews
Taitel 2014	Wrong intervention - pharmacy-led compliance intervention, not remote reviews
Uysal 2013	Wrong intervention - validating the Asthma Control Test via text messaging
van den Berg 2002	Wrong intervention - general practitioner (GP) telephone access to paediatricians
van der Meer 2009	Wrong intervention - technology-based self management between reviews
van Gaalen 2012	Wrong intervention - multifaceted intervention, not just remote reviews
van Reisen 2010	Wrong intervention - multifaceted intervention, not just remote reviews
Vasbinder 2013	Wrong intervention - minimal or no provider involvement. Medication reminder system
Vollmer 2006	Wrong intervention - technology-based self management between reviews
Voorend-van Bergen 2013	Wrong intervention - FeNO and Internet-based monitoring
Wiecha 2007	Wrong intervention - multi-faceted intervention, not just about remote monitoring
Willems 2008	Wrong intervention - technology-based self management between reviews
Young 2012	Wrong intervention - technology-based self management between reviews

(Continued)

Yun 2013	Wrong intervention - asthma education via text, not remote reviews
Zachgo 2002	Wrong intervention - computer works out best inhaler type for patient

Abbreviations: RCT = randomised controlled trial; PEF = peak expiratory flow.

DATA AND ANALYSES

Comparison 1. Remote versus face-to-face asthma reviews

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Exacerbations requiring oral corticosteroids	2		Odds Ratio (M-H, Random, 95% CI)	Totals not selected
1.1 Efficacy randomised controlled trials (RCTs)	1		Odds Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.2 Cluster implementation study	1		Odds Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2 Exacerbations requiring hospital emergency department (ED) treatment	4		Odds Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Efficacy RCTs	3	651	Odds Ratio (M-H, Random, 95% CI)	2.60 [0.63, 10.64]
2.2 Cluster implementation study	1	1212	Odds Ratio (M-H, Random, 95% CI)	1.19 [0.38, 3.71]
3 Exacerbations requiring hospital admission	4		Peto Odds Ratio (Peto, Fixed, 95% CI)	Subtotals only
3.1 Efficacy RCTs	3	651	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.63 [0.06, 6.32]
3.2 Cluster implementation study	1	1213	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.18 [0.83, 5.69]
4 Asthma control (Asthma Control Questionnaire (ACQ))	2		Mean Difference (IV, Random, 95% CI)	Totals not selected
4.1 Efficacy RCTs	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
4.2 Cluster implementation study	1		Mean Difference (IV, Random, 95% CI)	0.0 [0.0, 0.0]
5 Serious adverse events (including mortality)	1		Odds Ratio (M-H, Random, 95% CI)	Totals not selected
5.1 Cluster implementation study	1		Odds Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
6 Asthma-related quality of life (Asthma Quality of Life Questionnaire (AQLQ))	4		Mean Difference (IV, Random, 95% CI)	Subtotals only
6.1 Efficacy RCTs	3	544	Mean Difference (IV, Random, 95% CI)	0.08 [-0.14, 0.30]
6.2 Cluster implementation study	1	536	Mean Difference (IV, Random, 95% CI)	-0.02 [-0.23, 0.19]
7 Unscheduled healthcare visits	3		Odds Ratio (M-H, Random, 95% CI)	Subtotals only
7.1 Efficacy RCTs	2	531	Odds Ratio (M-H, Random, 95% CI)	0.91 [0.45, 1.85]
7.2 Cluster implementation study	1	1213	Odds Ratio (M-H, Random, 95% CI)	0.95 [0.75, 1.21]
8 Change in lung function (trough FEV ₁)	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
9 Adverse events	1	278	Odds Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

Comparison 2. Remote versus face-to-face for OCS tapering


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Exacerbations requiring hospital admission	1		Odds Ratio (M-H, Random, 95% CI)	Totals not selected
2 Asthma control (ACQ)	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
3 Asthma-related quality of life (AQLQ)	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
4 Unscheduled healthcare visits	1		Odds Ratio (M-H, Random, 95% CI)	Totals not selected
5 Adverse events	1		Odds Ratio (M-H, Random, 95% CI)	Totals not selected

Analysis 1.1. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 1 Exacerbations requiring oral corticosteroids.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 1 Exacerbations requiring oral corticosteroids




Analysis 1.2. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 2 Exacerbations requiring hospital emergency department (ED) treatment.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 2 Exacerbations requiring hospital emergency department (ED) treatment




Analysis 1.3. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 3 Exacerbations requiring hospital admission.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 3 Exacerbations requiring hospital admission




Analysis 1.4. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 4 Asthma control (Asthma Control Questionnaire (ACQ)).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 4 Asthma control (Asthma Control Questionnaire (ACQ))



(1) change from baseline, 0 to 12 months


(2) endpoint scores at 12 months

Analysis 1.5. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 5 Serious adverse events (including mortality).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 5 Serious adverse events (including mortality)




(1) Just mortality

Analysis 1.6. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 6 Asthma-related quality of life (Asthma Quality of Life Questionnaire (AQLQ)).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 6 Asthma-related quality of life (Asthma Quality of Life Questionnaire (AQLQ))



(1) Child scale data used. Parent scale is reported narratively.

(2) Endpoint scores at 12 months

(3) Endpoint scores - trial length variable


(4) Endpoint scores at 12 months

Analysis 1.7. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 7 Unscheduled healthcare visits.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 7 Unscheduled healthcare visits



(1) Not described as 'unscheduled' - total GP visits


(2) Face-to-face GP and specialist care control groups combined

Analysis 1.8. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 8 Change in lung function (trough FEV₁).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 8 Change in lung function (trough FEV₁)




(1) Face-to-face GP and specialist care control groups combined

Analysis 1.9. Comparison 1 Remote versus face-to-face asthma reviews, Outcome 9 Adverse events.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 1 Remote versus face-to-face asthma reviews

Outcome: 9 Adverse events




Analysis 2.1. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 1 Exacerbations requiring hospital admission.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 2 Remote versus face-to-face for OCS tapering

Outcome: 1 Exacerbations requiring hospital admission




Analysis 2.2. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 2 Asthma control (ACQ).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 2 Remote versus face-to-face for OCS tapering

Outcome: 2 Asthma control (ACQ)




Analysis 2.3. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 3 Asthma-related quality of life (AQLQ).

Review: Remote versus face-to-face check-ups for asthma

Comparison: 2 Remote versus face-to-face for OCS tapering

Outcome: 3 Asthma-related quality of life (AQLQ)




Analysis 2.4. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 4 Unscheduled healthcare visits.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 2 Remote versus face-to-face for OCS tapering

Outcome: 4 Unscheduled healthcare visits




Analysis 2.5. Comparison 2 Remote versus face-to-face for OCS tapering, Outcome 5 Adverse events.

Review: Remote versus face-to-face check-ups for asthma

Comparison: 2 Remote versus face-to-face for OCS tapering

Outcome: 5 Adverse events



ADDITIONAL TABLES

Table 1. Summary of study and intervention characteristics

Study ID	Total N	Country	Duration	Mean age	% male	% FEV ₁	Intervention	Control
Chan 2007	120	Hawaii, USA	12 months	9.6	62.5	100.5	In-home, website-based case management and education.	In-person education and case management.
Gruffydd-Jones 2005	194	UK	12 months	50.2	45.4	NR	6-monthly phone calls from trained asthma nurses. Formulation of individual AAP	6-monthly usual face-to-face appointment with an asthma nurse. Symptoms, peak flow and inhaler technique checked, and participants were issued with an AAP
Hashimoto 2011	95	The Netherlands	6 months	50.1	45.3	73.9	OCS dose adjustment guided by an internet-based diary, decision support and monitoring with support from a study nurse	OCS dose adjustment according to GINA by the specialist.

Table 1. Summary of study and intervention characteristics (Continued)

Pinnock 2003	278	UK	3 months	55.5	41.4	NR	Telephone check-up with the asthma nurse.	Face-to-face check-ups in the surgery with the asthma nurse.
Pinnock 2007a	1728	UK	12 months	42.6	44.6	NR	Three invitations to book either a telephone or face-to-face check-up. Non-attenders were phoned and reviewed opportunistically	Three invitations to book a face-to-face check-up. Non-attenders were not phoned opportunistically
Rasmussen 2005	300	Denmark	6 months	29	34.5	92.0	Participants were given an AAP, on-line electronic diary and peak flow meter. Physicians gave participants instructions via e-mail or telephone aided by computer decision support	The specialists taught the participants how to adjust their medication on the basis of a peak flow meter and AAP

Total N: the total number of participants randomised in the study, included to groups not analysed in this Cochrane review

% FEV₁: the baseline mean of the predicted normal values

Abbreviations: AAP = asthma action plan; NR = not reported; OCS = oral corticosteroids

APPENDICES

Appendix I. Sources and search methods for the Cochrane Airways Review Group's Specialised Register (CAGR)

Electronic searches: core databases

Database	Search frequency
CENTRAL (the Cochrane Library)	Monthly
MEDLINE (Ovid)	Weekly
Embase (Ovid)	Weekly
PsycINFO (Ovid)	Monthly
CINAHL (EBSCO)	Monthly
AMED (EBSCO)	Monthly

Hand-searches: core respiratory conference abstracts

Conference	Years searched
American Academy of Allergy, Asthma and Immunology (AAAAI)	2001 onwards
American Thoracic Society (ATS)	2001 onwards
Asia Pacific Society of Respirology (APSR)	2004 onwards
British Thoracic Society Winter Meeting (BTS)	2000 onwards
Chest Meeting	2003 onwards
European Respiratory Society (ERS)	1992, 1994, 2000 onwards
International Primary Care Respiratory Group Congress (IPCRG)	2002 onwards
Thoracic Society of Australia and New Zealand (TSANZ)	1999 onwards

MEDLINE search strategy used to identify trials for the CAGR

Asthma search

1. exp Asthma/
2. asthma\$.mp.
3. (antiasthma\$ or anti-asthma\$).mp.
4. Respiratory Sounds/
5. wheez\$.mp.

6. Bronchial Spasm/
7. bronchospas\$.mp.
8. (bronch\$ adj3 spasm\$).mp.
9. bronchoconstrict\$.mp.
10. exp Bronchoconstriction/
11. (bronch\$ adj3 constrict\$).mp.
12. Bronchial Hyperreactivity/
13. Respiratory Hypersensitivity/
14. ((bronchial\$ or respiratory or airway\$ or lung\$) adj3 (hypersensitiv\$ or hyperreactiv\$ or allerg\$ or insufficiency)).mp.
15. ((dust or mite\$) adj3 (allerg\$ or hypersensitiv\$)).mp.
16. or/1-15

Filter to identify randomised controlled trials (RCTs)

1. exp "clinical trial [publication type]" /
2. (randomised or randomised).ab,ti.
3. placebo.ab,ti.
4. dt.fs.
5. randomly.ab,ti.
6. trial.ab,ti.
7. groups.ab,ti.
8. or/1-7
9. Animals/
10. Humans/
11. 9 not (9 and 10)
12. 8 not 11

The MEDLINE strategy and RCT filter are adapted to identify trials in other electronic databases.

Appendix 2. Search strategy to identify relevant trials from the CAGR

- #1 AST:MISC1
- #2 MeSH DESCRIPTOR Asthma Explode All
- #3 asthma*:ti,ab
- #4 #1 or #2 or #3
- #5 MeSH DESCRIPTOR Telemedicine Explode All
- #6 telehealth* or tele-health*
- #7 telemedicine* or tele-medicine*
- #8 (internet* or computer* or web*):ti,ab,kw
- #9 interactive* or telecommunication*
- #10 (telephone or phone or SMS):ti,ab,kw
- #11 tele-monitor* or telemonitor*
- #12 telemanagement or tele-management
- #13 teleconsultation or tele-consultation
- #14 telecare* or tele-care*
- #15 telematic*
- #16 telepharmacy or tele-pharmacy
- #17 telenurs* or tele-nurs*
- #18 (video or email or e-mail):ti,ab,kw
- #19 remote NEXT consult*
- #20 wireless or bluetooth
- #21 tele-homecare or telehomecare
- #22 "remote care"
- #23 tele-support or telesupport

#24 mobile NEXT health*

#25 "computer mediated therapy"

#26 ehealth or e-health

#27 mhealth or m-health

#28 #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27

#29 #4 and #28

[Note: in search line #1, MISC1 denotes the field in which the reference has been coded for condition, in this case, asthma]

CONTRIBUTIONS OF AUTHORS

Kayleigh Kew (KK) wrote the text of the [Background](#) and [Methods](#), with significant comments and clinical input from Christopher Cates (CJC). Both review authors extracted and checked the data. KK contacted study authors for additional data, entered data into the analyses and wrote up the results. Both review authors contributed to the results interpretation, grading of the evidence and preparation of the final manuscript.

DECLARATIONS OF INTEREST

Kayleigh Kew has no known conflicts of interest.

Christopher Cates has no known conflicts of interest.

SOURCES OF SUPPORT

Internal sources

- Kayleigh Kew, UK.

Supported by St George's, University of London

External sources

- National Institute for Health Research, UK.

Evidence to guide care in adults and children with asthma, 13/89/14

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We assessed participant and personnel blinding separately for the objective and subjective outcomes, which we had not planned in the protocol ([Kew 2015b](#)). We were unable to conduct the subgroup analyses for age and type of technology due to an insufficient number of included studies.

We included exacerbations that required hospital admission rather than adverse events in the 'Summary of findings' table. We could not include both as we had to keep to seven outcomes to adhere to guidelines, and we considered the hospital admission data to be more important than all adverse events which tended to be reported as part of the exacerbation and resource use data in the studies.

We removed a sentence about searching manufacturer websites from the methods as it came from a template and is irrelevant to this research question.

As discussed in the 'Unit of analysis issues' section, we did not anticipate the inclusion of a cluster randomised controlled trial (RCT) so we had not outlined how we would deal with [Pinnock 2007a](#), a large two-cluster implementation study that we identified. We included it because it met the other inclusion criteria, but we presented it separately from the other studies due to the differences in

the study's design and analyses. For clarity in the analyses and write-up, we referred to [Pinnock 2007a](#) as the 'cluster implementation study' and [Chan 2007](#), [Gruffydd-Jones 2005](#), [Pinnock 2003](#) and [Rasmussen 2005](#) as the 'efficacy RCTs'. There were only two clusters so we included the data with participants as the unit of analysis.