



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Data on prevalence and risk factors associated with *Toxocara* spp infection, atopy and asthma development in Northeast Brazilian school children



Márcia B. Silva^a, Ana L.M. Amor^b, Leonardo N. Santos^a,
Alana A. Galvão^a, Aida Y. Oviedo V^a, Eduardo S. Silva^a,
Cynara Gomes Barbosa^c, Philip J. Cooper^{d,e},
Camila A. Figueiredo^a, Rita de Cassia Ribeiro^f,
Neuza Maria Alcântara-Neves^{a,*}

^a Instituto de Ciências da Saúde, Universidade Federal da Bahia, Salvador, Bahia, Brazil

^b Centro de Ciências da Saúde. Universidade Federal do Recôncavo da Bahia, Santo Antônio de Jesus, Bahia, Brazil

^c Faculdade de Farmácia, Universidade Federal da Bahia, Salvador, Bahia, Brazil

^d Facultad de Ciencias Médicas, de la Salud y la Vida, Universidad Internacional del Ecuador, Quito, Ecuador

^e St George's University of London, Division of Clinical Sciences, Cranmer Terrace, London SW17 0RE, UK

^f Escola de Nutrição, Universidade Federal da Bahia, Salvador, Brazil

ARTICLE INFO

Article history:

Received 12 April 2016

Received in revised form

11 August 2016

Accepted 22 August 2016

Available online 17 September 2016

Keywords:

Risk factors

ABSTRACT

In the present article, we provide shortly, data on risk factors for acquiring *Toxocara* spp. infection and investigate possible associations between this infection with atopy and asthma in school children of a small town and its semi-rural areas of Northeast Brazil. The data set are composed by demographic, social and home environment variables. The detection of anti-*Toxocara* spp. IgG and specific IgE to aeroallergens was determined by ELISA and ImmunoCAP/Phadiatrop systems, respectively. The data presented in this article are related to the article entitled "Risk factors for

DOI of original article: <http://dx.doi.org/10.1016/j.actatropica.2016.04.005>

* Correspondence to: Laboratório de Alergia e Acarologia, Instituto de Ciências da Saúde, Universidade Federal da Bahia, Avenida Reitor Miguel Calmon, sem nº, Canela, CEP: 40110-100, Salvador, Bahia, Brazil.

E-mail addresses: marciasilva25j@yahoo.com.br (M.B. Silva), ana_moreno22@yahoo.com.br (A.L.M. Amor), nsleo@yahoo.com.br (L.N. Santos), alana_alcantara@hotmail.com (A.A. Galvão), goviv1@yahoo.com (A.Y. Oviedo V), eduardossilva06@gmail.com (E.S. Silva), cynara.barbosa@gmail.com (C.G. Barbosa), pcooper@sgul.ac.uk (P.J. Cooper), cavfigueiredo@gmail.com (C.A. Figueiredo), ritaribeiroufba@gmail.com (R.d.C. Ribeiro), neuzalcantara@gmail.com (N.M. Alcântara-Neves).

<http://dx.doi.org/10.1016/j.dib.2016.08.062>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Toxocara spp, Atopy
Wheezing/Asthma

Toxocara spp. seroprevalence and its association with atopy and asthma phenotypes in school-age children in a small town and semi-rural areas of Northeast Brazil” (M.B. Silva, A.L. Amor, L.N. Santos, A.A. Galvão, A.V. Oviedo Vera, E.S. Silva et al., 2016) [1].

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Epidemiology, allergology
More specific subject area	Immunoparasitology
Type of data	Figure, Table
How data was acquired	ELISA, ImmunoCAP and Phadiatropo
Data format	Analyzed
Experimental factors	Stool and blood samples, and sera for measurement of allergen-specific IgE and anti- <i>Toxocara</i> spp. IgG
Experimental features	Immunoassay
Data source location	Federal University of Bahia, Salvador, Bahia, Brazil
Data accessibility	Data is available with this article

Value of the data

- These data set will be of value for the scientific community who work in the area of infectious diseases since it involves the risk factors related to *Toxocara* spp. infection.
- The data will also be of value for studies in the area of allergy and its interface with helminthic diseases, since they report an association of *Toxocara* spp. infection with aeroallergen specific IgE.
- These data reinforce the hypothesis that this association may be related to the cross-reactivity between parasite-specific and aeroallergen-specific IgE.

1. Data

The data demonstrate obtained in this work is summarized in Fig. 1 and Table 1. Fig. 1 shows that being male and having contact with dogs and cats were risk factors for *Toxocara* spp. infection among other variables studied. Table 1 shows the analysis of *Toxocara* spp. infection as risk factors for atopy and asthma. We found that *Toxocara* spp. seropositive school age children were more prone to have positive serum aeroallergen-specific IgE.

2. Experimental design, materials and methods

The data presented in this paper investigated the risk factors to acquire *Toxocara* spp. infection and its association with atopy and asthma [1]. It reports the data obtained in a study conducted in a small city of Northeast Brazil with 791 school-age children. Students underwent an epidemiological survey answered by their parents on sanitation, social class and risk factors for toxocariasis. Moreover, they answered an ISAAC Portuguese-adapted questionnaire for asthma diagnosis [2]. Blood collection was

Associations between potential risk factors and IgG seropositivity to *Toxocara* spp.




Fig. 1. ORs and 95% confident intervals calculated by logistic analysis model adjusted by age, sex, contact with dogs and cats, school location, maternal schooling, family income and helminth infection.

Table 1

Association between *Toxocara* spp. seropositivity with atopy and wheezing/asthma in 791 elementary school students, 6–13 years old.

Outcomes	n (%) / N	Adjusted OR (95% CI) ^a
Phadiatop* IgE ≥ 0.70 KU/L	251(49.9)/503	1.95 (1.40–2.72)
<i>B. tropicalis</i> specific IgE ≥ 0.70 KU/L	286 (56.8)/503	1.85 (1.31–2.62)
Any allergen IgE ≥ 0.70 KU/L	342 (67.9)/503	2.00 (1.49–2.68)
Atopic wheezing/asthma	398(79.2)/503	1.04 (0.54–2.08)
Non-atopic wheezing/asthma	57(11.3)/503	1.08 (0.40–2.70)

^a IgE specific to *Blomia tropicalis* (D201) and to Phadiatop aeroallergens (pollen extracts, fungi extracts, dog and cat epithelia and *Dermatophagoides* spp.) measured by immunoCAP.




Fig. 2. Schematic representation of the steps involved during the study to achieve the results.

performed to evaluate the levels of allergen-specific IgE and IgG anti-*Toxocara* spp. reactivity. Univariable and multivariable analyses were used to analyze *Toxocara* spp. infection risk factors and the association of this infection with atopy and asthma phenotypes. The variables investigated were: gender, age, maternal schooling, income, school location, contact with dog and cat (Fig. 2).

Conflict of interest

The authors declare that they have no competing interests.

Acknowledgments

The authors would like to acknowledge FAPESB, Brazil, for financial support and technical support to conduct this work.

Transparency document. Supplementary material

Transparency data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.08.062>.

References

- [1] M.B. Silva, A.L. Amor, L.N. Santos, A.A. Galvão, A.V. Oviedo Vera, E.S. Silva, et al., Risk factors for *Toxocara* spp. seroprevalence and its association with atopy and asthma phenotypes in school-age children in a small town and semi-rural areas of Northeast Brazil, *Acta Trop.* (2016).
- [2] A. Strina, S. Cairncross, M.L. Barreto, C. Larrea, M.S. Prado, Childhood diarrhea and observed hygiene behavior in Salvador, Brazil, *Am. J. Epidemiol.* (2003).