

SUPPLEMENTARY TABLES & FIGURES:


SUPPLEMENTARY TABLE 1: INFORMATION ON THE DERIVATION AND EXTERNAL VALIDATION STUDIES

Study [ref]	Date of Study	Location	Setting	Age Group	Response Rate	Ethnicity Reporting
Derivation Datasets:						
1) The assessment of Body Composition in Children (ABCC) Study [17]	2011-2012	London	Primary schools	8-10 years	64%	Reported by parent
2) The East London Bioelectrical Impedance (ELBI) [23]	2008-2009	London	Secondary schools, weight management clinic and volunteers recruited by advertisement	11-15 years	Not Applicable	Self-reported
3) The Reference Child (RC) [24]	2001-2010	London	Volunteers recruited by advertisement	4-22 years	Not Applicable	Self-reported or reported by parent
4) The Size and Lung function in Children (SLIC) Study [25]	2012-2013	London	Primary Schools	5-11 years	52%	Reported by parent
Validation Dataset:						
Avon Longitudinal Study of Parents and Children (ALSPAC) [26, 27]	2002-2003	Bristol	Population based advertising	11-12 years	Not Applicable	Reported by parent

SUPPLEMENTARY TABLE 2: ASSESSMENT OF CALIBRATION SLOPE AND CALIBRATION-IN-THE-LARGE IN TERMS OF FAT MASS AND FAT FREE MASS FROM INTERNAL-EXTERNAL CROSS-VALIDATION


Study omitted for external validation	Fat Free Mass		Fat Mass	
	Calibration-In-The-Large	Calibration Slope	Calibration-In-The-Large	Calibration Slope
1	0.05 (-0.45 to 0.55)	0.97 (0.95 to 0.99)	0.35 (0.16 to 0.54)	1.03 (1.01 to 1.05)
2	1.62 (0.34 to 2.90)	0.98 (0.95 to 1.01)	-0.73 (-1.37 to -0.10)	0.99 (0.95 to 1.03)
3	1.02 (0.36 to 1.67)	0.98 (0.96 to 1.01)	-0.74 (-1.15 to -0.33)	1.02 (0.98 to 1.05)
4	-0.37 (-0.97 to 0.22)	1.05 (1.02 to 1.07)	-0.20 (-0.48 to 0.09)	0.95 (0.93 to 0.97)
Pooled	0.46 (-0.30 to 1.21)	1.00 (0.96 to 1.03)	-0.29 (-0.83 to 0.25)	1.00 (0.95 to 1.04)

SUPPLEMENTARY FIGURE 1: HISTOGRAM OF FAT FREE MASS (TOP) AND LN(FAT FREE MASS), BY SEX




FOOTNOTE: LN = Natural logarithmic transformation

SUPPLEMENTARY FIGURE 2: CALIBRATION SLOPES AND CALIBRATION-IN-THE-LARGE (KG), BY SEX, FROM INTERNAL-EXTERNAL CROSS VALIDATION



FOOTNOTE: Study codes: 1 = The assessment of Body Composition in Children Study, 2 = The East London Bioelectrical Impedance, 3 = The Reference Child, 4 = The Size and Lung function in Children Study.

SUPPLEMENTARY FIGURE 3: CALIBRATION SLOPES AND CALIBRATION-IN-THE-LARGE (KG), BY ETHNIC GROUP, FROM INTERNAL-EXTERNAL CROSS VALIDATION



FOOTNOTE: Study codes: 1 = The assessment of Body Composition in Children Study, 2 = The East London Bioelectrical Impedance, 3 = The Reference Child, 4 = The Size and Lung function in Children Study. Ethnic group labels: 1 = White, 2 = Black, 3 = South Asian, 4 = Other Asian, 5 = Other.

APPENDIX 1: COMPARISON OF ESTIMATING FM DIRECTLY OR INDIRECTLY

The two approaches for estimating FM can be written as follows:

Equation 1: $FM_{\text{direct}} = f(\text{height, weight, age, sex, ethnic group}) + \varepsilon$; such that $\varepsilon \sim N(0, \sigma^2)$

Equation 2: $FM_{\text{indirect}} = \text{weight} - \text{FFM} = \text{weight} - [g(\text{height, weight, age, sex, ethnic group}) + \mu]$; such that $\mu \sim N(0, \tau^2)$.

Therefore, possible estimates of FM are \hat{f} or $\text{weight} - \hat{g}$. It was clear from scatter plots of both FM and FFM, obtained from the DD method, plotted against height (Figure 1 below), that the variance in FFM with height was more homogeneous than for FM (i.e. $\tau^2 < \sigma^2$). Hence, a regression model for FFM (equation 2) should be estimated with greater precision than a model for FM (equation 1) because of the more homogenous relationship with height. As the variance of weight is ≈ 0 due to negligible measurement error, equation 2 ($\text{weight} - \hat{g}$) would be expected to provide more precise estimates of FM compared to estimates from \hat{f} , making the indirect approach likely to be the preferred method for estimating FM.




Figure 1: Scatterplot of FM (top) and FFM (bottom) against height in boys (left) and girls (right)

APPENDIX 2: INTERNAL VALIDATION VIA BOOTSTRAPPING

The following steps were undertaken for the bootstrapping process³³:

1. 1000 bootstrap samples were randomly selected from the entire DD derivation datasets, with replacement, such that the size of each bootstrap sample ($N=2375$) was equal to that of the entire DD derivation dataset ($N=2375$). This selection process was stratified by sex, ethnic group and age to ensure that each bootstrap sample contained a representative sample of each of the subgroups.
2. The final developed prediction model was fitted within each bootstrap sample to obtain estimates of model performance based on R^2 , the calibration slope and calibration-in-the-large (CIL).
3. Calculation of optimism-adjustments for R^2 , the calibration slope and CIL:
 - a. Values of R^2 , the calibration slope and CIL were obtained, for each 1000 bootstrap sample
 - b. The values from a. were subtracted from the original R^2 , the calibration slope and CIL values from the original DD derivation datasets to obtain the level of optimism in each performance measure within each of the 1000 bootstrap samples
 - c. The average of the differences in measures from b. across the 1000 bootstrap samples were determined
 - d. Optimism-adjusted values of R^2 (denoted as R^2_{adjusted}), the calibration slope (denoted as Calibration Slope_{adjusted}) and CIL (denoted as CIL_{adjusted}) were obtained by subtracting the average of the differences (i.e. the value from c.) from the original R^2 and calibration slopes from the original DD derivation datasets